



Code: UKCAL-CWF-CON-EIA-RPT-00007-7D02

Volume 7D Caledonia South Appendices

Appendix 3-2 Sediment Quality Baseline

Caledonia Offshore Wind Farm Ltd

5th Floor Atria One, 144 Morrison Street, Edinburgh, EH3 8EX



Volume 7D Appendix 3-2 Sediment Quality Baseline

| | |
|-----------------|----------------------------------|
| Code | UKCAL-CWF-CON-EIA-RPT-00007-7D02 |
| Revision | Issued |
| Date | 18 October 2024 |

Table of Contents

| | | |
|-----|---------------------------------|----|
| 1 | Introduction | 1 |
| 2 | Sediment Characterisation | 3 |
| 2.1 | Overview | 3 |
| 2.2 | Caledonia South Site..... | 5 |
| 2.3 | Caledonia South OECC | 8 |
| 3 | Total Organic Carbon..... | 12 |
| 3.1 | Overview | 12 |
| 3.2 | Caledonia South Site..... | 12 |
| 3.3 | Caledonia South OECC | 12 |
| 4 | References | 13 |

List of Figures

| | |
|--|---|
| Figure 1-1: Site-Specific Survey Locations in the Caledonia South Site and Caledonia South OECC..... | 2 |
| Figure 2-1: Existing surficial sediment characterisation combined with site-specific sampling..... | 4 |

List of Tables

| | |
|---|---|
| Table 2-1: Summary of Particle Size Analysis across the Caledonia South Site. | 6 |
| Table 2-2: Summary of Particle Size Analysis across the Caledonia South OECC. | 9 |

Acronyms and Abbreviations


| | |
|------------------|-----------------------------------|
| BGS | British Geological Survey |
| EIA | Environmental Impact Assessment |
| MW&SQ | Marine Water and Sediment Quality |
| OECC | Offshore Export Cable Corridor |
| OWF | Offshore Wind Farm |
| PSA | Particle Size Analysis |
| TOC | Total Organic Carbon |

1 Introduction

1.1.1.1 This appendix covers an in-depth analysis of the sediment quality baseline environment of Caledonia South in terms of sediment quality parameters in the Marine Water and Sediment Quality (MW&SQ) Study Area, such as:

- Sediment Characterisation (including Particle Size Analysis (PSA)); and
- Total Organic Carbon (TOC) content.

1.1.1.2 The location of sample stations across the Caledonia South Site and the Caledonia South Offshore Export Cable Corridor (OECC) are shown in Figure 1-1. The results of this site-specific survey are compared with publicly available regional data, including previous Offshore Wind Farm (OWF) Environmental Impact Assessments (EIAs), to provide a robust characterisation of the entire MW&SQ Study Area.



- Caledonia South Site
- Caledonia South Offshore Export Cable Corridor
- 10km Zone of Influence
- PSA Sample
- Chemistry Sample
- × Water Sample

Service Layer Credits: © OpenStreetMap (and) contributors, CC-BY-SA
 Esri, Garmin, GEBCO, NOAA NGDC, and other contributors
 © Caledonia Offshore Wind Farm Ltd © 2024. This document is the property of contractors and sub-contractors and shall not be reproduced nor transmitted without prior written approval.





| | | | |
|--|--|-------------------------------------|--|
| | | | |
| | | | |
| <small>CONTRACTOR DRAWING NO</small> UKCAL_1_GO_WNF_MWS_MAP_00439 | | <small>CONTRACTOR REV</small> 01 | |
| <small>COORDINATE PARAMETERS</small> WGS 84 / UTM zone 30N (EPSG: 32630) | | | |
| <small>DRAWING TITLE</small> Figure 1-1: Site-Specific Survey Locations in the Caledonia South Site and Caledonia South OECC | | | |
| <small>STATUS</small> Approved | | <small>SCALE</small> 1:500,000 | |
| <small>DRAWING NUMBER</small> N/A | | <small>SHEET NO</small> 01 of 01 | |
| | | <small>REV</small> N/A | |

2 Sediment Characterisation

2.1 Overview

2.1.1.1 Sediment characteristics (grain type and size) influence multiple of important biotic and abiotic aspects in the marine environment and are therefore considered an important impact pathway. For example, diversity and composition of macro and meio benthic communities, organic carbon storage and contaminant storage/distribution are strongly conditioned by sediment grain size. Sediments with a finer particle size, such as clays and muds, can act as adsorption surfaces for contaminants that may be released into the water column if the sediment is disturbed. This is due to their larger surface area to volume ratios and higher organic carbon content. Sediments consisting of coarser sand and gravel are accepted to carry a much lower contamination risk. Information regarding particle sizes is an important step in assessing the contamination risk to the marine environment. Figure 2-1 illustrates the anticipated sediment characterisation across the MW&SQ Study Area.



- Caledonia South Site
- Caledonia South Offshore
- Export Cable Corridor
- 10km Zone of Influence



Seabed Sediments (Folk, 1954)

- 1.1.1 Mud
- 1.2.1 sandy Mud
- 1.2.2 (gravelly) sandy Mud
- 1.3.1 muddy Sand
- 1.3.2 (gravelly) muddy Sand
- 2.1.1 Sand
- 2.1.2 (gravelly) Sand
- 3.1.1 gravelly Sand
- 3.2.1 sandy Gravel
- 3.3.1 Gravel
- 4.3.1 gravelly muddy Sand
- 4.4.1 muddy sandy Gravel
- 5. Rock and Boulders

PSA Results (%)

- Fines
- Sand
- Gravel

Service Layer Credits: © OpenStreetMap (and) contributors, CC-BY-SA
 © Caledonia Offshore Wind Farm Ltd © 2024. This document is the property of contractors and sub-contractors and shall not be reproduced nor transmitted without prior written approval.



| REV | DATE | DOC STATUS | ORGIN | REVIEW | APP |
|-----|------------|------------|-------|--------|-----|
| 01 | 16/09/2024 | Approved | EV | BB | DH |

CONTRACTOR DRAWING NO: UKCAL1_CO_WNF_MWS_MAP_00440
 CONTRACTOR REV: 01
 GEOGRAPHIC PARAMETERS: WGS 84 / UTM zone 30N (EPSG: 32630)

DRAWING TITLE: **Figure 2-1: Existing Surficial Sediment Characterisation Combined with Site-Specific Sampling**

| | |
|---------------------|--------------------|
| STATUS: Approved | SCALE: 1:500,000 |
| DRAWING NUMBER: N/A | SHEET NO: 01 of 01 |
| | REV: N/A |

2.2 Caledonia South Site

- 2.2.1.1 Regional-scale sediment maps sourced from EMODnet (2020¹), derived from BGS datasets (Figure 2-1), indicate that within the Caledonia South Site, the surficial seabed sediments predominantly consist of Holocene sediments, including sand, gravelly sand, sandy gravel, and slightly gravelly sand, as per the Folk classification (Folk, 1954²). These sediments are reported to contain only a small proportion of fines, less than 10%. It is important to acknowledge that the BGS sediment maps are constructed from a relatively sparse sampling density in this area. Nevertheless, these data can serve to provide broad-scale characterization, particularly when combined with site-specific sampling campaigns.
- 2.2.1.2 A site-specific survey (Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)) sampled 17 stations across the Caledonia South Site using wet and dry sieving to determine the PSA of sediment grab samples, results of which are shown in Table 2-1 and Figure 2-1. The mean particle size ranged from 45µm at Station ENV21 to 1057µm at Station ENV22, with an average mean grain diameter of 253µm (±261SD). The dominant fraction, accounting for between 58.1% and 93.8% of the sediment was sand (with a particle size range of >63µm and <2mm). These data generally support the BGS sediment maps (i.e., reporting of less than 10% fines), where there was a slight exceedance of average portion of fines present in the survey samples of 13.9%. Surveys conducted across Moray East Offshore Wind Farm found levels of fine (silt/clay) particles were generally low across the survey array (<3%) but increased (around 4% and 5%) with increased water depth to the south (Moray Offshore Renewables Limited, 2012³). This is further validated by comparison with surveys undertaken by Moray West Offshore Wind Farm that found seabed sediments across the Moray West site generally consist of Holocene gravelly sand and sand with only a minor proportion of fines (<5 to 10% silt and clay sized) (Moray Offshore Windfarm (West) Limited, 2018⁴).
- 2.2.1.3 When survey results were classified under the modified Folk 16 classification (Folk, 1954²), stations ranged from muddy sand to gravelly sand. These classifications were validated by the analysis of the relative proportions of fines, sand and gravel. Similar results as seen reported from Moray East survey samples are observed in the Caledonia South Site where samples indicate a particle size distribution ranging from very poorly sorted to moderately sorted (Folk and Ward, 1957⁵). "Very poorly sorted" indicates that the sediment contains a wide range of particle sizes, with no dominant size fraction. This suggests a lack of uniformity in the particle sizes present. "Moderately sorted" suggests that there is some degree of uniformity in the particle sizes, but there is still noticeable variability present.

Table 2-1: Summary of Particle Size Analysis across the Caledonia South Site.

| Station | Mean Diameter (µm) | Fines (%) | Sand (%) | Gravel (%) | Sorting | Modified Folk Classification | Total Organic Carbon (%) * |
|---------|--------------------|-----------|----------|------------|-----------|------------------------------|----------------------------|
| ENV05 | 160 | 14.1 | 85.7 | 0.1 | Poor | Muddy sand | N/A |
| ENV06 | 208 | 9.3 | 90.4 | 0.3 | Poor | Sand | N/A |
| ENV10 | 763 | 2.8 | 86.2 | 11.0 | Poor | Gravelly sand | N/A |
| ENV12 | 106 | 18.2 | 81.8 | 0.0 | Poor | Muddy sand | 0.44 |
| ENV15 | 157 | 11.0 | 87.9 | 1.1 | Poor | Slightly gravelly muddy sand | 0.44 |
| ENV16 | 125 | 21.1 | 78.3 | 0.5 | Poor | Muddy sand | N/A |
| ENV19 | 182 | 14.4 | 85.6 | 0.0 | Poor | Muddy sand | N/A |
| ENV21 | 45 | 41.9 | 58.1 | 0.0 | Very poor | Muddy sand | 1.10 |
| ENV22 | 1057 | 8.5 | 68.0 | 23.5 | Very poor | Gravelly muddy sand | 0.43 |
| ENV24 | 197 | 8.9 | 90.7 | 0.4 | Poor | Sand | 0.32 |
| ENV25 | 229 | 16.4 | 77.4 | 6.2 | Very poor | Gravelly muddy sand | N/A |
| ENV26 | 120 | 15.6 | 84.4 | 0.0 | Poor | Muddy sand | N/A |
| ENV27 | 325 | 4.5 | 84.3 | 11.2 | Poor | Gravelly sand | N/A |
| ENV29 | 190 | 8.4 | 91.0 | 0.6 | Poor | Sand | N/A |

| Station | Mean Diameter (µm) | Fines (%) | Sand (%) | Gravel (%) | Sorting | Modified Folk Classification | Total Organic Carbon (%) * |
|---|--------------------|-----------|----------|------------|-----------------------|------------------------------|----------------------------|
| ENV30 | 85 | 21.6 | 78.4 | 0.0 | Poor | Muddy sand | N/A |
| ENV32 | 230 | 5.5 | 93.8 | 0.7 | Moderate | Sand | N/A |
| ENV34 | 124 | 14.4 | 85.5 | 0.1 | Poor | Muddy sand | 0.45 |
| Summary Statistics | | | | | | | |
| Minimum | 45 | 2.8 | 58.1 | 0.0 | Very poor to Moderate | Muddy sand to Gravelly sand | 0.32 |
| Maximum | 1057 | 41.9 | 93.8 | 23.5 | Very poor to Moderate | Muddy sand to Gravelly sand | 1.10 |
| Mean | 253 | 13.9 | 82.8 | 3.3 | Very poor to Moderate | Muddy sand to Gravelly sand | 0.53 |
| Standard Deviation | 261 | 9.1 | 9.1 | 6.4 | Very poor to Moderate | Muddy sand to Gravelly sand | 0.28 |
| * TOC was analysed via chemistry samples acquired at nine sample stations. Therefore, N/A indicates where TOC was not analysed. | | | | | | | |

2.3 Caledonia South OECC

- 2.3.1.1 Regional-scale sediment maps from EMODnet (2020¹), sourced from BGS datasets, indicate that within the Caledonia South OECC the surficial seabed sediments are characterised by Holocene sediments of sand and gravels close to the Caledonia South Site, with mud content increasing towards the shore (Figure 2-1). Sediments generally appear to become progressively finer as water depth increases, with isolated patches of coarser sediment associated with bathymetric highs. Between approximately 8 and 20km offshore, the surficial sediments are classified as muddy sand, with a band of sandy mud approximately 10km offshore (Figure 2-1).
- 2.3.1.2 Across the Caledonia South OECC, 38 stations were sampled using wet and dry sieving to determine the PSA of sediment grab samples as part of a site-specific survey (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)). Results, detailed in Table 2-2 and Figure 2-1, revealed a wide range in mean particle size, ranging from 41µm (Station ENV27) to 1813µm (Station ENV11). Stations closest to the shore, situated in water depths less than 35m (ENV07, ENV53, ENV57, ENV58, and ENV59), were predominantly sandy (>95%) and exhibited moderate to moderately well sorted characteristics based on Folk and Ward (1957⁵) statistics. These stations recorded minimal fines (<5%) and gravel. In contrast, stations in water depths exceeding 70m (ENV02 to ENV06, ENV08 to ENV10, ENV12, ENV17, ENV19, ENV23 to ENV31, ENV33 to ENV36) displayed poorly to very poorly sorted sediment, classified as muddy sand under modified Folk classification (1954²). These stations were primarily composed of fine sand to very fine sand, with fines content ranging from 11% at Station ENV08 to 45% at Station ENV27, and gravel content below 1%. Overall, there was a clear positive correlation between fines and water depth, where finer sediment was observed in areas of deeper water.

Table 2-2: Summary of Particle Size Analysis across the Caledonia South OECC.

| Station | Mean Diameter (µm) | Fines (%) | Sand (%) | Gravel (%) | Sorting | Modified Folk Classification | Total Organic Carbon (%) * |
|---------|--------------------|-----------|----------|------------|-----------|------------------------------|----------------------------|
| ENV01 | 1591 | 1.8 | 59.6 | 38.6 | Very poor | Sandy gravel | N/A |
| ENV02 | 90 | 27.3 | 72.6 | <0.1 | Very poor | Muddy sand | N/A |
| ENV03 | 115 | 19.1 | 80.9 | 0.0 | Poor | Muddy sand | N/A |
| ENV04 | 61 | 31.3 | 68.7 | 0.0 | Poor | Muddy sand | 0.37 |
| ENV05 | 83 | 22.1 | 77.9 | <0.1 | Poor | Muddy sand | N/A |
| ENV06 | 141 | 14.1 | 85.9 | <0.1 | Poor | Muddy sand | N/A |
| ENV07 | 520 | 1.7 | 97.4 | 0.8 | Moderate | Sand | N/A |
| ENV08 | 167 | 11.1 | 88.9 | <0.1 | Poor | Muddy sand | N/A |
| ENV09 | 79 | 30.5 | 69.5 | 0.0 | Very poor | Muddy sand | N/A |
| ENV10 | 51 | 34.9 | 65.1 | 0.0 | Poor | Muddy sand | N/A |
| ENV11 | 1813 | 6.2 | 56.2 | 37.5 | Very poor | Sandy gravel | N/A |
| ENV12 | 69 | 30.6 | 69.4 | 0.0 | Very poor | Muddy sand | N/A |
| ENV13 | 220 | 9.5 | 90.3 | 0.2 | Poor | Sand | 0.19 |
| ENV14 | 215 | 16 | 81.4 | 2.6 | Poor | Slightly gravelly muddy sand | N/A |
| ENV15 | 490 | 6.1 | 79.8 | 14.1 | Poor | Gravelly sand | N/A |

| Station | Mean Diameter (µm) | Fines (%) | Sand (%) | Gravel (%) | Sorting | Modified Folk Classification | Total Organic Carbon (%) * |
|---------|--------------------|-----------|----------|------------|-----------|------------------------------|----------------------------|
| ENV16 | 363 | 4.9 | 92.4 | 2.7 | Poor | Slightly gravelly sand | N/A |
| ENV17 | 130 | 18.9 | 80.8 | 0.3 | Poor | Muddy sand | 0.30 |
| ENV18 | 166 | 24.6 | 64.7 | 10.7 | Very poor | Gravelly muddy sand | N/A |
| ENV19 | 118 | 19.1 | 80.6 | 0.3 | Poor | Muddy sand | N/A |
| ENV21 | 497 | 8.7 | 72.1 | 19.1 | Very poor | Gravelly muddy sand | N/A |
| ENV22 | 252 | 12.3 | 87.2 | 0.5 | Poor | Muddy sand | N/A |
| ENV23 | 152 | 16.2 | 83.8 | 0.0 | Poor | Muddy sand | N/A |
| ENV24 | 53 | 37.1 | 62.9 | 0.0 | Very poor | Muddy sand | N/A |
| ENV25 | 143 | 12.2 | 87.8 | <0.1 | Poor | Muddy sand | N/A |
| ENV26 | 47 | 40.7 | 59.3 | 0.0 | Very poor | Muddy sand | 0.77 |
| ENV27 | 41 | 45.2 | 54.8 | 0.0 | Very poor | Muddy sand | N/A |
| ENV28 | 66 | 29.7 | 70.3 | 0.0 | Poor | Muddy sand | N/A |
| ENV29 | 48 | 37.2 | 62.8 | 0.0 | Poor | Muddy sand | 0.61 |
| ENV30 | 91 | 20.4 | 79.6 | <0.1 | Poor | Muddy sand | N/A |
| ENV31 | 99 | 23.2 | 76.8 | 0.0 | Poor | Muddy sand | 0.35 |
| ENV33 | 111 | 18.7 | 81.3 | 0.0 | Poor | Muddy sand | N/A |

| Station | Mean Diameter (µm) | Fines (%) | Sand (%) | Gravel (%) | Sorting | Modified Folk Classification | Total Organic Carbon (%) * |
|--|--------------------|-----------|----------|------------|------------------------------|------------------------------|----------------------------|
| ENV34 | 73 | 25.3 | 74.6 | 0.1 | Poor | Muddy sand | N/A |
| ENV35 | 102 | 24.4 | 75.6 | <0.1 | Very poor | Muddy sand | N/A |
| ENV36 | 107 | 16 | 84 | <0.1 | Poor | Muddy sand | 0.62 |
| ENV53 | 270 | 2.5 | 95.1 | 2.4 | Moderately well | Slightly gravelly sand | 0.17 |
| ENV57 | 194 | 1.7 | 98.3 | <0.1 | Moderately well | Sand | N/A |
| ENV58 | 228 | 1.7 | 98 | 0.3 | Moderately well | Sand | 0.16 |
| ENV59 | 350 | 1.1 | 98.8 | 0.1 | Moderately well | Sand | N/A |
| Summary Statistics | | | | | | | |
| Minimum | 41 | 1.1 | 54.8 | 0.0 | Very poor to moderately well | Muddy sand to sandy gravel | 0.16 |
| Maximum | 1813 | 45.2 | 98.8 | 38.6 | Very poor to moderately well | Muddy sand to sandy gravel | 0.77 |
| Mean | 248 | 18.5 | 78 | 3.4 | Very poor to moderately well | Muddy sand to sandy gravel | 0.40 |
| Standard Deviation | 371 | 12.2 | 12.3 | 9.2 | Very poor to moderately well | Muddy sand to sandy gravel | 0.43 |
| * TOC was analysed via chemistry samples acquired at nine sample stations. Therefore, N/A indicates where TOC was not analysed | | | | | | | |

3 Total Organic Carbon

3.1 Overview

3.1.1.1 TOC is expressed as a percentage of the total weight and represents the carbon constituent of organic matter. The primary components of organic matter in marine sediment are detrital matter, naphthenic material (e.g., carboxylic acids) and humic substances with a small proportion of biological biomass. In general, for continental shelf sediment there is a close relationship between organic carbon content and the surface area of the mineral matrix (Mayer, 1994⁶). Increased TOC is expected with the presence of fine sediment, as it adsorbs to the increased surface area provided by the fine grain particles.

3.2 Caledonia South Site

3.2.1.1 A site-specific survey found that across the Caledonia South Site, TOC concentrations ranged from 0.23% at Station ENV07 to 1.10% at ENV21, where the mean TOC value was 0.49%. Station ENV21 was identified as a high outlier, also reporting the highest fines content. In the Caledonia South Site, these TOC values were expected considering the general predominant sand component of the sediment (see Table 2-1). This was supported by results where TOC concentration was positively correlated with the volume of fines across the Caledonia South Site.

3.3 Caledonia South OECC

3.3.1.1 A site-specific survey found that across the Caledonia South OECC, TOC concentrations ranged from 0.16% at Station ENV58 to 0.77% at Station ENV26, where the mean TOC value was 0.39% and there was a positive correlation with water depth. Similar to the Caledonia South Site, these TOC values were expected in the Caledonia South OECC considering the general predominant sand and muddy sand components of the sediment (see Table 2-2). This was supported by results where TOC concentration was positively correlated with the volume of fines across the Caledonia South OECC.

4 References

- ¹ EMODnet (2020) 'European Marine Observation and Data Network (EMODnet) Bathymetry data'. Available at: <https://portal.emodnet-bathymetry> (Accessed 01/05/2024)
- ² Folk, R.L. (1954) 'The distinction between grain size and mineral composition in sedimentary-rock nomenclature'. *The Journal of Geology* 62(4): 344-359
- ³ Moray Offshore Renewables Limited (2012) 'Moray East OWF Environmental Statement (Volume 2, Chapter 4: Biological Environment Baseline)'. Available at: https://marine.gov.scot/sites/default/files/chapter_4_-_biological_environment_baseline.pdf (Accessed 30/05/2024)
- ⁴ Moray Offshore Windfarm (West) Limited (2018) 'Moray West OWF Offshore EIA Report (Technical Appendix 7.1: Benthic Survey Report)'. Available at: https://marine.gov.scot/sites/default/files/00538036_3.pdf (Accessed 30/05/2024)
- ⁵ Folk, R.L. and Ward, W.C. (1957) 'Brazos River Bar: A study in the significance of grain size parameters'. *Journal of Sedimentary Petrology* 27(1): 3-26
- ⁶ Mayer, L.M. (1994) 'Surface area control on organic carbon accumulation in continental shelf sediments'. *Geochimica et Cosmochimica Acta* 58: 1271-84

Caledonia Offshore Wind Farm
5th Floor, Atria One
144 Morrison Street
Edinburgh
EH3 8EX

www.caledoniaoffshorewind.com

