



**Code:** UKCAL-CWF-CON-EIA-RPT-00007-7C03

# **Volume 7C Caledonia North Appendices**

## Appendix 3-3 Sediment Contamination Baseline

Caledonia Offshore Wind Farm Ltd

5th Floor Atria One, 144 Morrison Street, Edinburgh, EH3 8EX



# Volume 7C Appendix 3-3 Sediment Contamination Baseline

<b>Code</b>	UKCAL-CWF-CON-EIA-RPT-00007-7C03
<b>Revision</b>	Issued
<b>Date</b>	18 October 2024

# Table of Contents

1	Introduction.....	1
2	Sediment Contamination Assessment Guidelines .....	3
2.1	Action Levels .....	3
2.2	Canadian Marine Sediment Quality Guidelines .....	5
2.3	OSPAR Assessment Criteria .....	6
3	Sediment Contamination Baseline .....	8
4	Total Hydrocarbon Content .....	11
4.1	Overview .....	11
4.2	Caledonia North Site .....	11
4.3	Caledonia North OECC.....	14
5	Polycyclic Aromatic Hydrocarbons.....	17
5.1	Overview .....	17
5.2	Caledonia North Site .....	17
5.3	Caledonia North OECC.....	20
6	Polychlorinated Biphenyls .....	23
6.1	Overview .....	23
6.2	Caledonia North Site .....	23
6.3	Caledonia North OECC.....	23
7	Polybrominated Diphenyl Ethers.....	24
7.1	Overview .....	24
7.2	Caledonia North Site .....	24
7.3	Caledonia North OECC.....	25
8	Organochloropesticides .....	26
8.1	Overview .....	26
8.2	Caledonia North Site .....	26
8.3	Caledonia North OECC.....	26
9	Organotins .....	27
9.1	Overview .....	27
9.2	Caledonia North Site .....	27
9.3	Caledonia North OECC.....	27
10	Metals .....	28

10.1	Overview.....	28
10.2	Caledonia North Site .....	28
10.3	Caledonia North OECC .....	29
11	Sediment Contamination Conclusions .....	31
12	References.....	32

## List of Figures

Figure 1-1: Site-specific survey locations in the Caledonia North Site and Caledonia North OECC. ....2

## List of Tables

Table 2-1: Action Levels used in sediment contaminant assessment as presented in Marine Scotland (2017). .....	4
Table 2-2: Canadian Marine Sediment Quality Guidelines for PAHs (CCME, 1999). .....	6
Table 2-3: OSPAR Assessment Criteria for Polybrominated Diphenyl Ethers (OSPARa, 2020; OSPARb, 2020; Vinas <i>et al.</i> , 2023). .....	7
Table 4-1: Summary of sediment hydrocarbon analysis of samples collected within the Caledonia North Site. ....	13
Table 4-2: Summary of sediment hydrocarbon analysis of samples collected within the Caledonia North OECC.....	15
Table 5-1: Summary of PAH concentration from sediment samples across the Caledonia North Site. ....	18
Table 5-2: Summary of PAH concentrations from sediment samples across the Caledonia North OECC. ....	21
Table 7-1: Concentration of PBDE 209 identified when normalised to 2.5% TOC from sediment samples across the Caledonia North Site. ....	24
Table 7-2: Concentration of PBDE 209 identified when normalised to 2.5% TOC from sediment samples across the Caledonia North OECC.....	25
Table 10-1: Summary statistics of sediment metal contamination from samples collected across the Caledonia North Site. ....	29
Table 10-2: Summary statistics of sediment metal contamination from samples collected across the Caledonia North OECC. ....	30
Table 11-1 Summary of sediment contamination in the Caledonia North Site and Caledonia North OECC relative to threshold guidelines. ....	31

## Acronyms and Abbreviations

<b>AL</b>	Action Level
<b>BAC</b>	Background Assessment Concentration
<b>BC</b>	Background Concentration
<b>CCME</b>	Canadian Council of Ministers of the Environment
<b>CPI</b>	Carbon Preference Index
<b>CSEMP</b>	Clean Seas Environment Monitoring Programme
<b>EAC</b>	Environmental Assessment Criteria
<b>EIA</b>	Environmental Impact Assessment
<b>EQS</b>	Environmental Quality Standards
<b>EQSD</b>	Environmental Quality Standards Directive
<b>ERL</b>	Effects Range Low
<b>ERM</b>	Effects Range Median
<b>FEQG</b>	Federal Environmental Quality Guidelines
<b>GC</b>	Gas Chromatography
<b>GC-FID</b>	Gas Chromatography with Flame Ionization Detection
<b>HMW</b>	High Molecular Weight
<b>LMW</b>	Low Molecular Weight
<b>LOD</b>	Limit of Detection
<b>MMO</b>	Marine Management Organisation
<b>MW&amp;SQ</b>	Marine Water and Sediment Quality
<b>OCP</b>	Organochloropesticide
<b>OECC</b>	Offshore Export Cable Corridor


<b>OSPAR</b>	The Convention for the Protection of the Marine Environment of the North-East Atlantic
<b>OWF</b>	Offshore Wind Farm
<b>PAH</b>	Polycyclic Aromatic Hydrocarbon
<b>PBDE</b>	Polybrominated Diphenyl Ether
<b>PCB</b>	Polychlorinated Biphenyl
<b>PEL</b>	Probable Effect Level
<b>SD</b>	Standard Deviation
<b>TEL</b>	Threshold Effect Level
<b>THC</b>	Total Hydrocarbon Content
<b>TOC</b>	Total Organic Carbon
<b>TBT</b>	Tributyltin
<b>UCM</b>	Unresolved Complex Mixture
<b>USEPA</b>	United States Environmental Protection Agency
<b>UVF</b>	Ultraviolet Fluorescence
<b>WFD</b>	Water Framework Directive

# 1 Introduction

1.1.1.1 This appendix provides a detailed analysis of the baseline environment of Caledonia North in terms of sediment contamination in the Marine Water and Sediment Quality (MW&SQ) Study Area, including:


- Sediment contamination assessment guidelines;
- Total Hydrocarbon Content (THC);
- Polycyclic Aromatic Hydrocarbons (PAHs);
- Polychlorinated Biphenyls (PCBs);
- Polybrominated Diphenyl Ethers (PBDEs);
- Organochloropesticides (OCPs);
- Organotin; and
- Metals.

1.1.1.2 Illustrations of the location of sample stations across the Caledonia North Site (i.e., Array Area) and Caledonia North Offshore Export Cable Corridor (OECC) are shown in Figure 1-1. The results of this site-specific survey are compared with publicly available regional data, including previous Offshore Wind Farm (OWF) Environmental Impact Assessments (EIAs) and the OSPAR Intermediate Assessment, to provide a robust characterisation of the entire MW&SQ Study Area.



- Caledonia North Site
- Caledonia North Offshore Export Cable Corridor
- 10km Zone of Influence
- PSA Sample
- Chemistry Sample
- × Water Sample

Service Layer Credits: © OpenStreetMap (and) contributors, CC-BY-SA  
 Esri, Garmin, GEBCO, NOAA NGDC, and other contributors  
 © Caledonia Offshore Wind Farm Ltd © 2024. This document is the property of contractors and sub-contractors and shall not be reproduced nor transmitted without prior written approval.



REV	DATE	DOC STATUS	ORIGIN	REVIEW	APP	
01	16/09/2024	Approved		EV	BB	DH

CONTRACTOR DRAWING NO UKCAL1_GO_WNF_MWS_MAP_00437	CONTRACTOR REV 01
GEOGRAPHIC PARAMETERS WGS 84 / UTM zone 30N (EPSG: 32630)	
Figure 1-1: Site-Specific Survey Locations in the Caledonia North Site and Caledonia North OECC	

STATUS Approved	SCALE 1:500,000
DRAWING NUMBER N/A	SHEET NO 01 of 01
	REV N/A

## 2 Sediment Contamination Assessment Guidelines

### 2.1 Action Levels

- 2.1.1.1 In the absence of Environmental Quality Standards (EQSs) for sediments, survey sediment contaminant data has been analysed relative to Action Levels (ALs) for the disposal of dredged material. Following the Marine Directorate pre-disposal sampling guidance (Marine Scotland, 2017<sup>1</sup>; note, in 2023, Marine Scotland was renamed Marine Directorate), the assessment of material should be based on these ALs, which are presented in Table 2-1. In the current assessment, these ALs are used to determine whether further assessment is required.
- 2.1.1.2 Contaminants below Action Level 1 (AL1) are, for dredging projects, not considered to be of concern and thus can be disposed of at sea. Contaminant levels which exceed Action Level 2 (AL2) are not considered suitable for disposal at sea. Those sediments which record concentrations between AL1 and AL2 may be disposed of at sea, but may require some consideration prior to doing so.
- 2.1.1.3 Whilst Caledonia North is not a dredging project per se, it does involve a proposal to dredge, drill and dispose (i.e., disturbance) of seabed sediments within the Caledonia North Site and Caledonia North OECC. Therefore, adhering to guidelines regarding the disposal of dredged material will ensure a cautious assessment of potential environmental impacts (IMO, 2014<sup>2</sup>; OSPAR, 2014<sup>3</sup>).

Table 2-1: Action Levels used in sediment contaminant assessment as presented in Marine Scotland (2017).

Contaminant	Action Level 1 (mg/kg; Dry Weight) (ppm)	Action Level 2 (mg/kg; Dry Weight) (ppm)
Arsenic	20	70
Cadmium	0.4	4
Chromium	50	370
Copper	30	300
Lead	50	400
Mercury	0.25	1.5
Nickel	30	150
Zinc	130	600
Organotins (e.g., Tributyltin (TBT))	0.1	0.5
Polychlorinated Biphenyls (PCBs) – sum of 25 congeners	0.02	0.18
Polycyclic Aromatic Hydrocarbons (PAHs)*	0.1	-
Total Hydrocarbon Content (THC)	100	-
* Dibenzo(a,h)anthracene (PAH) has an AL1 of 0.01mg/kg dry weight (ppm) (no AL2).		

2.1.1.4 The standard procedure in reviewing PAH concentrations in marine sediment samples is to consider against the Effects Range Low (ERL) and the Effects Range Median (ERM) for a discrete suite of low molecular weight (LMW) and high molecular weight (HMW) PAHs (Gorham-Test *et al.*, 1999<sup>4</sup>). This effectively presents a similar AL1 (ERL) and AL2 (ERM) approach to provide context to sediment quality for PAHs and has been applied to support this MW&SQ assessment. The sum of the following PAH concentrations is used in the calculations:

- HMW PAHs: Fluoranthene, Pyrene, Benz[a]anthracene, Chrysene, Benzo(a)pyrene, Dibenz(a,h)anthracene; and
- LMW PAHs: Naphthalene, Acenaphthene, Fluorene, Anthracene, C1-naphthalenes, Acenaphthylene, Phenanthrene.

- 2.1.1.5 The ERL (equivalent to AL1) for the sum of LMW and HMW PAHs is 552 and 1,700µg/kg, respectively. The ERM (equivalent to AL2) for the sum of LMW and HMW PAHs is 3,160 and 9,600µg/kg, respectively.

## 2.2 Canadian Marine Sediment Quality Guidelines

- 2.2.1.1 The Canadian Marine Sediment Quality Guidelines (Table 2-2) have also been used within this assessment to provide context to the sediment contaminant levels reported within the site-specific samples. These Guidelines provide some information for those contaminants not currently reported within the ALs (Table 2-1), specifically PAHs. Developed by the Canadian Council of Ministers of the Environment (CCME), these Guidelines are applied within this Project to provide an indication on the degree of sedimentary contamination and the likely ecological impact.

- 2.2.1.2 The Guidelines allow the identification of three ranges of chemical contaminants, with regard to biological effects:

- Below the Threshold Effect Levels (TEL): the minimal effect range within which adverse effects rarely occur;
- Between the TEL and Probable Effect Levels (PEL): the possible effect range within which adverse effects occasionally occur; and
- Above the PEL: the probable effect range within which adverse effects frequently occur.

Table 2-2: Canadian Marine Sediment Quality Guidelines for PAHs (CCME, 1999<sup>5</sup>).

Contaminant	Threshold Effect Levels (µg/kg)	Probable Effect Levels (µg/kg)
Acenaphthene	6.71	88.9
Acenaphthylene	5.87	128
Anthracene	46.9	245
Benz(a)anthracene	74.8	693
Benzo(a)pyrene	88.8	763
Chrysene	108	846
Dibenz(a,h)anthracene	6.22	135
Fluoranthene	113	1,494
Fluorene	21.2	144
2-Methylnaphthalene	20.2	201
Naphthalene	34.6	391
Phenanthrene	86.7	544
Pyrene	153	1,398

## 2.3 OSPAR Assessment Criteria

2.3.1.1 In the absence of ALs for Polybrominated Diphenyl Ethers (PBDEs) in sediment, levels used by OSPAR (OSPAR, 2023<sup>6</sup>), presented in Table 2-3, have been adopted here:

- Background Assessment Criteria (BAC); and
- Federal Environmental Quality Guidelines (FEQGs) also referred to as Environmental Assessment Criteria (EAC).

Table 2-3: OSPAR Assessment Criteria for Polybrominated Diphenyl Ethers (OSPARa, 2020<sup>7</sup>; OSPARb, 2020<sup>8</sup>; Vinas *et al.*, 2023<sup>9</sup>).

PBDE Congener	BAC (mg/kg)	FEQG (mg/kg)
BDE28	0.00005	0.11
BDE47	0.00005	0.0975
BDE66	0.00005	0.0975
BDE85	0.00005	0.001
BDE99	0.00005	0.001
BDE100	0.00005	0.001
BDE126	0.00005	n/a
BDE153	0.00005	1.1
BDE154	0.00005	1.1
BDE183	0.00005	14
BDE209	0.00005	0.0475

## 3 Sediment Contamination Baseline

3.1.1.1 Elevated levels of contaminants can affect flora and fauna in a variety of ways, ranging from cellular effects in individuals to ecosystem effects resulting from changes in population sizes or even the loss of an entire species (UK Marine SACs Project, 2001<sup>10</sup>). Therefore, it is important to accurately investigate the potential sediment contamination present in the receiving environment of Caledonia North to ensure we understand the full potential of impacts. Unlike water quality, there are no formal quantitative EQS for sediments. Therefore, in the absence of any quantified UK standards, common practice for characterising baseline sediment quality conditions is to compare against AIs, as is suggested for the disposal of dredged material (Marine Scotland, 2017<sup>1</sup>). Moreover, where ALs do not provide comprehensive guidance, other sediment quality guidelines are used in place (See Section 2).

3.1.1.2 The Intermediate Assessment 2017 prepared by OSPAR (2017<sup>11</sup>) reviewed and compared mercury, cadmium, lead, organotin (limited data), PCB, PAH and PBDE concentrations in sediments between OSPAR contaminant assessment areas (including data collated as part of the Clean Seas Environmental Monitoring Programme (CSEMP) assessment). Four monitoring stations in the Moray Firth form part of the Northern North Sea region. The main conclusions drawn from the data for the Northern North Sea are as follows:

- Metals:
  - Mercury and lead concentrations in sediment are at or above the BAC.
  - Lead concentrations are also at or above the ERL value.
  - Mean concentrations of cadmium are below the BAC.
- PCBs:
  - Trend analysis shows PCB concentrations are reducing.
  - PCB congener 118 concentrations in sediments are below the EAC and above the BAC.
- PAHs:
  - Mean PAH concentrations in sediment were statistically significantly below the ERL but not below the BAC.
- PBDEs:
  - PBDE concentrations in sediment are low and often below detection levels.

3.1.1.3 Further to the OSPAR (2017<sup>11</sup>) assessment, there is already a comprehensive base understanding of the volume and type of contamination likely to be present around the MW&SQ Study Area,

particularly due to the previous Environmental Impact Assessment Reports (EIARs) reports carried out for preceding OWFs built in the surrounding area.

- 3.1.1.4 In the EIA conducted for the most recently consented OWF in the Moray Firth, Moray West, detailed sediment chemistry samples were examined (Moray West Offshore Wind Farm Ltd, 2018<sup>12</sup>). The analysis of heavy metals in the subtidal samples revealed concentrations well below the respective guidelines, with no samples exceeding ALs or the Canadian Sediment Quality Guidelines. Additionally, PAH concentrations were consistently low, often falling below the limit of detection (LOD) for the analytical tests. However, it's worth noting that the LODs for Acenaphthene, Acenaphthylene, and Dibenzo(ah)anthracene were slightly higher than the Canadian TEL values. Overall, the survey findings led to the conclusion that there were no indications of unacceptable or significant contamination levels in the vicinity.
- 3.1.1.5 In the EIA conducted for Moray East, sediment samples were analysed for metals (arsenic, cadmium, chromium, copper, mercury, nickel, lead, tin, barium, aluminium), PAHs (16 US EPA Priority Pollutants), total PAH and total petroleum hydrocarbons. Results showed concentrations were below respective guideline values (ALs and Canadian Sediment Quality Guidelines) (Moray Offshore Renewables Limited, 2012a<sup>13</sup>). No adverse environmental effects were anticipated and not considered further in the assessment (Moray Offshore Renewables Limited, 2012b<sup>14</sup>).
- 3.1.1.6 In the EIA conducted for Beatrice OWF, also within the Moray Firth, the survey findings closely mirror those of Moray West, despite the historical presence of oil and gas exploration and production activities near Beatrice (BOWL, 2012<sup>15</sup>). Organic contaminants (PAHs, n-alkanes, and total alkanes) were present at very low levels or not detected. No heavy metals were present at levels that exceeded ALs. There was no apparent relationship between distribution of any of the metals, including barium, and the distribution of known wellheads (barium compounds, such as barite (barium sulfate), are commonly used as weighting agents in drilling fluids for oil and gas wells). Barium was present at levels ranging from 133 to 273 mg/kg, and whilst this may have come from past drilling activities, these levels are lower than published values for North Sea sediments sampled at greater than 5km from oil and gas platforms, and do not suggest gross contamination. This assumption is further supported by a recent decommissioning plan for the Beatrice Field in the Central North Sea. Contaminant concentrations varied throughout the survey area; elevated concentrations of contaminants were found in the Beatrice Alpha Drilling and Accommodation Platform cuttings pile beneath the platform, and within the surrounding area within 100 m, whilst samples from the wider Beatrice area showed the concentrations of contaminants were generally within expected background levels for the Central North Sea.

Further indicating a lack of migrating contamination from these oil and gas areas (Repsol Sinopec Resources UK Limited, 2018<sup>16</sup>).

#### 3.1.1.7

Upon comparing the above surveys carried out for the application of previous OWFs, it becomes evident that sediment contamination estimates provided by OSPAR (2017<sup>11</sup>) appear slightly conservative for this specific area, where less prominent sediment contamination has so far been observed. The following sections discuss, in detail, the types of contamination and the site-specific surveys of the Caledonia North Site (Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)) and Caledonia North OECC (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)).

## 4 Total Hydrocarbon Content

### 4.1 Overview

- 4.1.1.1 The principal sources of hydrocarbons in the marine environment are anthropogenic (McDougall, 2000<sup>17</sup>). However, biosynthetic, and geochemical processes can also be a source for hydrocarbons in the marine environment. THC concentration gives an indication of the total hydrocarbon present within a sediment sample. In the case of site-specific survey output, THC is equivalent to total n-alkane (nC10 to nC37), pristane, phytane, Unresolved Complex Mixture (UCM) and total PAH (all PAHs including alkylated derivatives) concentrations.
- 4.1.1.2 It has previously been shown that benthic macrofauna suffer adverse effects when THC is in excess of 50 µg/g (Kjeilen-Eilertsen *et al.*, 2004<sup>18</sup>; UKOOA, 2002<sup>19</sup>; 2005<sup>20</sup>) and as such, this value represents a threshold above which hydrocarbons can be expected to present a 'significant environmental impact'. In addition, Kingston (1992) previously reported that benthic macrofauna suffer adverse effects (e.g. reduced diversity), when THC is more than 50µg/g to 60µg/g, where specific sensitive species can experience impacts at levels greater than 10µg/g (Kingston, 1992<sup>21</sup>).
- 4.1.1.3 THC concentrations give an indication of the total oil in the sediment at each station and can therefore be compared to the ecotoxicological limits outlined above. Analysis of GC chromatograms can be used to identify the source of hydrocarbons and the extent to which they are weathered. In particular, abundant UCM is ascribed to either degraded or weathered oil residues, and therefore its occurrence in environmental samples is an indicator of oil pollution (Bouloubassi *et al.*, 2001<sup>22</sup>). Notably, a UCM between nC20 and nC34, centred on nC29 is typical of North Sea sediments, and is generally considered as 'North Sea Background'.

### 4.2 Caledonia North Site

- 4.2.1.1 Analysis has been conducted using Gas Chromatography with Flame Ionization Detection (GC-FID) to measure the concentration of THC in sediment samples collected from different stations across the Caledonia North Site (Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)). The results showed THC concentration ranging from 2.1µg/g at Station ENV35 to 3.6µg/g at Station ENV07, with an average concentration of 3.0µg/g (± 0.65 SD). Overall, concentrations of THC by ultraviolet fluorescence (UVF) were lower than THC concentrations by GC-FID, where no Stations recorded concentrations above the LOD (Table 4-1).
- 4.2.1.2 The chromatograms presented a general pattern of low-level, high molecular weight (HMW) n-alkanes and UCM ranging from nC20 to nC36.

These distributions are considered typical of background levels of hydrocarbon inputs within the North Sea with historical oil and gas explorations which include a relatively low level UCM distributed between nC20 and nC33.

- 4.2.1.3 Hydrocarbons in the molecular weight range nC24 to nC36 commonly originate from terrestrial plant sources (Harborne, 1999<sup>23</sup>) or may represent the residue of highly weathered and biodegraded petrogenic material including natural seeps, shipping discharge and oil and gas exploration and extraction (Bouloubassi *et al.*, 2001<sup>22</sup>; McDougall, 2000<sup>17</sup>). The results of the THC profile is characteristic of background concentrations of n-alkanes in North Sea sediments, and it is indicative of sedimentary biogenic hydrocarbons from a terrestrial plant source (Tran *et al.*, 1995<sup>24</sup>). The UCM accounted for 43% to 79% of THC across the Caledonia North Site, indicating that generally hydrocarbons were weathered with some fresher inputs.
- 4.2.1.4 THC values across the Caledonia North Site were below published threshold values. Overall, the faunal community is not expected to be influenced by THC concentrations.

Table 4-1: Summary of sediment hydrocarbon analysis of samples collected within the Caledonia North Site.

Sample Station	THC by UVF	THC by GC-FID	UCM by GC-FID	n-alkanes by GC-FID			CPI by GC-FID	Isoprenoids by GC-FID		
				nC10-20	nC21-37	nC10-37		Pristane	Phytane	Pr/Ph Ratio
ENV07	<1	3.6	1.6	0.004	0.055	0.059	1.9	<1	<1	NC
ENV18	<1	3.1	2.0	0.010	0.033	0.044	2.4	0.002	<1	NC
ENV33	<1	3.3	2.1	0.015	0.036	0.051	3.4	0.003	<1	NC
ENV35	<1	2.1	1.2	0.001	0.015	0.016	NC	0.002	<1	NC
<b>Summary Statistics</b>										
Minimum	NQ	2.1	1.2	0.001	0.015	0.016	NQ	NQ	NQ	NC
Maximum	NQ	3.6	2.1	0.015	0.055	0.059	3.4	0.003	NQ	NC
Mean	NC	3.0	1.7	0.008	0.035	0.043	NC	NC	NC	NC
Standard Deviation	NC	0.65	0.4	0.006	0.016	0.019	NC	NC	NC	NC
Concentrations expressed as µg/g dry sediment. NQ: Not quantified due to concentrations below LOD. NC: Not calculated due to concentrations below LOD.										

## 4.3 Caledonia North OECC

- 4.3.1.1 Analysis has been conducted using GC-FID to measure the concentration of THC in sediment samples collected from different stations across the Caledonia North OECC (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)). The results showed THC concentration ranged from 2.8µg/g at the shallowest station (ENV58) up to 18.7µg/g at the deepest station (ENV26) with a mean of 10.7µg/g ( $\pm$  4.6 SD). The data suggest that concentrations of THC were positively correlated with fines and TOC. Similar to the Caledonia North Site, concentrations of THC by UVF were lower than THC concentrations by GC-FID, where only three Stations recorded concentrations above the LOD (Table 4-2).
- 4.3.1.2 Concentrations of THC by GC-FID at six of the nine stations (ENV04, ENV17, ENV26, ENV29 ENV36 and ENV53) were marginally above the 10µg/g threshold and as such have a potential for negative impacts on sensitive benthic macrofauna. Of note, concentrations of THC by UVF were below the threshold of 10µg/g.
- 4.3.1.3 The chromatograms generally presented a pattern comparable to samples collected in the Caledonia North Site of low-level, HMW n-alkanes and unresolved complex mixtures (UCM) ranging from nC20 to nC36. These distributions are considered typical of background levels of hydrocarbon inputs within the North Sea with historical oil and gas explorations which include a relatively low level UCM distributed between nC20 and nC33.
- 4.3.1.4 The results of the THC profile are characteristic of background concentrations of n-alkanes in North Sea sediments, and it is indicative of sedimentary biogenic hydrocarbons from a terrestrial plant source (Tran *et al.*, 1995<sup>24</sup>). The UCM accounted for 62% to 94% of THC across the Caledonia North Site, indicating that generally hydrocarbons were weathered with some fresher inputs.
- 4.3.1.5 Overall, THC concentrations along the Caledonia North OECC were generally below the published thresholds indicating the faunal community is not expected to be significantly influenced by THC concentrations.

Table 4-2: Summary of sediment hydrocarbon analysis of samples collected within the Caledonia North OECC.

Sample Station	THV by UVF	THC by GC-FID	UCM by GC-FID	n-alkanes by GC-FID			CPI by GC-FID	Isoprenoids by GC-FID		
				nC10-20	nC21-37	nC10-37		Pristane	Phytane	Pr/Ph Ratio
ENV04	<1	11.3	8.1	0.035	0.283	0.318	3.4	0.011	<0.001	NC
ENV13	<1	6.0	3.7	0.012	0.069	0.081	4.9	0.003	<0.001	NC
ENV17	4.1	12.9	8.2	0.033	0.331	0.364	3.7	0.010	<0.001	NC
ENV26	4.2	18.7	13.1	0.057	0.421	0.478	3.2	0.022	0.004	5.5
ENV29	4.7	14.0	9.5	0.047	0.314	0.361	3.0	0.012	0.003	4.4
ENV31	<1	9.2	6.4	0.027	0.175	0.202	3.6	0.007	0.002	4.0
ENV36	<1	11.4	7.4	0.040	0.211	0.251	4.2	0.010	0.002	5.7
ENV53	<1	10.1	9.5	0.003	0.016	0.018	NC	0.001	<0.001	NC
ENV58	<1	2.8	1.8	0.006	0.014	0.020	NC	0.001	<0.001	NC
<b>Summary Statistics</b>										
Minimum	NQ	2.8	1.8	0.003	0.014	0.018	NQ	0.001	NQ	NQ
Maximum	4.7	18.7	13.1	0.057	0.421	0.478	4.9	0.022	0.004	5.7

Sample Station	THV by UVF	THC by GC-FID	UCM by GC-FID	n-alkanes by GC-FID			CPI by GC-FID	Isoprenoids by GC-FID		
				nC10-20	nC21-37	nC10-37		Pristane	Phytane	Pr/Ph Ratio
Mean	NC	10.7	7.5	0.029	0.204	0.232	NC	0.009	NC	NC
±SD	NC	4.6	3.3	0.019	0.147	0.164	NC	0.006	NC	NC
Concentrations expressed as µg/g dry sediment. NQ: Not quantified due to concentrations below LOD. NC: Not calculated due to concentrations below LOD.										

## 5 Polycyclic Aromatic Hydrocarbons

### 5.1 Overview

5.1.1.1 PAHs are classed as priority hazardous substances and ubiquitous persistent, bio-accumulative and toxic compounds under the Water Framework Directive (WFD) in the related EQS Directive (2008/105/EC amended by 2013/39/EU). The Canadian Marine Sediment Quality Guidelines (TELs and PELs values) provide the most reliable standardised estimates of the potential toxicity of individual PAHs in marine sediments, as outlined in Section 2.2.

5.1.1.2 However, the concentrations at which individual PAHs produce toxic effects vary widely (Long *et al.*, 1995<sup>25</sup>) and are dependent on their type and bioavailability. Values for the toxicity of individual aromatics may be misleading since individual PAHs are rarely found in isolation. More sensitive estimates of the potential toxicity of PAHs in marine sediments are ERL and ERM concentrations for total LMW, total HMW and total PAHs. Long *et al.* (1995<sup>25</sup>) gives ERL concentrations for LMW and HMW PAHs of 0.55 and 1.70µg/g, respectively. ERM concentrations are 3.16 and 9.60 µg/g for LMW and HMW PAHs, respectively. The ERL and ERM concentrations for total PAH concentration in sediments are 4.022 and 44.792µg/g, respectively. These concentrations are not actual thresholds of toxicity but delineate concentration ranges with associated probabilities of toxicity.

### 5.2 Caledonia North Site

5.2.1.1 The total PAH concentrations ranged from 0.001µg/g to 0.004µg/g across the Caledonia North Site, except for Station ENV33 and ENV35 that were below even the LOD. The molecular mass indices identified the presence of pyrogenic PAHs (i.e., likely from grass, wood or coal combustion). Overall, concentrations of PAHs were very low, with no exceedance of either the Canadian or USEPA guidelines for PAH concentrations in sediment (shown in Table 5-1). Moreover, total PAH concentrations were below their respective ERL values (Long *et al.*, 1995<sup>25</sup>), further indicating that toxic effects to fauna by PAHs are unlikely.

Table 5-1: Summary of PAH concentration from sediment samples across the Caledonia North Site.

PAHs	Canadian Guidelines		Sample Station			
	TEL Value	PEL Value	ENV07	ENV18	ENV33	ENV35
Acenaphthene	6.71	88.9	<LOD	<LOD	<LOD	<LOD
Acenaphthylene	5.78	128	<LOD	<LOD	<LOD	<LOD
Anthracene	46.9	245	<LOD	<LOD	<LOD	<LOD
Benzo[a]anthracene	74.8	693	<LOD	<LOD	<LOD	<LOD
Benzo[a]pyrene	88.8	763	<LOD	<LOD	<LOD	<LOD
Benzo[b]fluoranthene	-	-	0.001	<LOD	<LOD	<LOD
Benzo[e]pyrene	-	-	0.001	<LOD	<LOD	<LOD
Benzo[ghi]perylene	-	-	<LOD	<LOD	<LOD	<LOD
Benzo[k]fluoranthene	-	-	<LOD	<LOD	<LOD	<LOD
C1-naphthalenes	-	-	<LOD	<LOD	<LOD	<LOD
C1-phenanthrenes	-	-	<LOD	<LOD	<LOD	<LOD
C2-naphthalenes	-	-	<LOD	<LOD	<LOD	<LOD
C3-naphthalenes	-	-	<LOD	<LOD	<LOD	<LOD

PAHs	Canadian Guidelines		Sample Station			
	TEL Value	PEL Value	ENV07	ENV18	ENV33	ENV35
Chrysene	108	846	<LOD	<LOD	<LOD	<LOD
Dibenzo[ah]anthracene	6.22	135	<LOD	<LOD	<LOD	<LOD
Fluoranthene	113	1,494	<LOD	<LOD	<LOD	<LOD
Fluorene	21.2	144	<LOD	<LOD	<LOD	<LOD
Indeno[1,2,3-cd]pyrene	-	-	0.002	0.001	<LOD	<LOD
Naphthalene	34.6	391	<LOD	<LOD	<LOD	<LOD
Perylene	-	-	<LOD	<LOD	<LOD	<LOD
Phenanthrene	86.7	544	<LOD	<LOD	<LOD	<LOD
Pyrene	153	1,398	<LOD	<LOD	<LOD	<LOD
Total	-	-	0.004	0.001	N/A	N/A
Concentrations from Sample Stations converted from ng/g dry weight sediment to ug/g dry weigh sediment to express comparisons with guideline levels.						

## 5.3 Caledonia North OECC

### 5.3.1.1

The total PAH concentrations ranged from 0.009µg/g to 0.146µg/g across the Caledonia North OECC, except for Station ENV53 and ENV58 that were below even the LOD. The molecular mass indices identified the presence of pyrogenic PAHs (i.e., likely from grass, wood or coal combustion) at all stations. Results indicate that PAHs were also from petrogenic (e.g., petroleum) in low concentrations. Overall, concentrations of PAHs were very low, with no exceedance of either the Canadian or USEPA guidelines for PAH concentrations in sediment (shown in Table 5-2). Moreover, total PAH concentrations were below their respective ERL values (Long *et al.*, 1995<sup>25</sup>), further indicating that toxic effects to fauna by PAHs are unlikely.

Table 5-2: Summary of PAH concentrations from sediment samples across the Caledonia North OECC.

PAHs	Canadian Guidelines		Sample Station								
	TEL Value	PEL Value	ENV04	ENV13	ENV17	ENV26	ENV29	ENV31	ENV36	ENV53	ENV58
Acenaphthene	6.71	88.9	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
Acenaphthylene	5.78	128	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
Anthracene	46.9	245	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
Benzo[a]anthracene	74.8	693	0.002	<LOD	0.002	0.005	0.003	0.002	0.002	<LOD	<LOD
Benzo[a]pyrene	88.8	763	0.003	<LOD	0.003	0.007	0.005	0.003	0.003	<LOD	<LOD
Benzo[b]fluoranthene	-	-	0.008	0.002	0.006	0.016	0.012	0.007	0.008	<LOD	<LOD
Benzo[e]pyrene	-	-	0.008	0.002	0.006	0.017	0.013	0.007	0.009	<LOD	<LOD
Benzo[ghi]perylene	-	-	0.005	<LOD	0.004	0.009	0.007	0.004	0.004	<LOD	<LOD
Benzo[k]fluoranthene	-	-	0.006	0.002	0.005	0.013	0.009	0.005	0.005	<LOD	<LOD
C1-naphthalenes	-	-	0.003	<LOD	0.002	<LOD	0.003	0.002	0.002	<LOD	<LOD
C1-phenanthrenes	-	-	0.003	<LOD	0.003	0.006	0.005	0.003	0.003	<LOD	<LOD
C2-naphthalenes	-	-	0.004	0.002	0.004	0.010	0.006	0.003	0.004	<LOD	<LOD
C3-naphthalenes	-	-	0.003	<LOD	0.002	0.005	0.004	0.002	0.002	<LOD	<LOD

PAHs	Canadian Guidelines		Sample Station								
	TEL Value	PEL Value	ENV04	ENV13	ENV17	ENV26	ENV29	ENV31	ENV36	ENV53	ENV58
Chrysene	108	846	0.003	<LOD	0.003	0.006	0.004	0.003	0.002	<LOD	<LOD
Dibenzo[ah]anthracene	6.22	135	<LOD	<LOD	<LOD	0.003	0.002	<LOD	<LOD	<LOD	<LOD
Fluoranthene	113	1,494	0.004	<LOD	0.004	0.009	0.006	0.005	0.003	<LOD	<LOD
Fluorene	21.2	144	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
Indeno[1,2,3-cd]pyrene	-	-	0.008	0.002	0.007	0.020	0.015	0.008	0.010	<LOD	<LOD
Naphthalene	34.6	391	<LOD	<LOD	<LOD	0.002	<LOD	<LOD	<LOD	<LOD	<LOD
Perylene	-	-	0.003	<LOD	0.004	0.007	0.002	<LOD	<LOD	<LOD	<LOD
Phenanthrene	86.7	544	0.002	<LOD	0.002	0.005	0.003	0.002	0.002	<LOD	<LOD
Pyrene	153	1,398	0.003	<LOD	0.003	0.007	0.005	0.004	0.003	<LOD	<LOD
Total	-	-	0.069	0.009	0.061	0.146	0.108	0.059	0.062	<LOD	<LOD
Concentrations from Sample Stations converted from ng/g dry weight sediment to ug/g dry weigh sediment to express comparisons with guideline levels.											

## 6 Polychlorinated Biphenyls

### 6.1 Overview

6.1.1.1 PCBs are synthetic organic compounds that are extremely harmful to the environment. PCBs are toxic and persistent compounds that are known to bioaccumulate in marine sediment, where they may be trapped in lower layers unless the sediments are disturbed. A report published by Marine Scotland (2020<sup>26</sup>) found PCB concentrations in biota (fish and shellfish) and sediment in the four Scottish biogeographic regions generally remain below the threshold at which adverse effects occur in marine life, but above Background Assessment Concentrations. Mean concentrations were stable or decreasing in the assessed areas over the assessment period (1999 – 2018) (Marine Scotland, 2020<sup>26</sup>).

### 6.2 Caledonia North Site

6.2.1.1 PCB concentrations from sediment samples collected from the Caledonia North Site were all below the LOD, indicating that toxic effects to fauna by PCBs are unlikely, even with the disruption of sediment.

### 6.3 Caledonia North OECC

6.3.1.1 PCB concentrations from sediment samples collected from the Caledonia North OECC were all below the LOD, indicating that toxic effects to fauna by PCBs are unlikely, even with the disruption of sediment.

## 7 Polybrominated Diphenyl Ethers

### 7.1 Overview

7.1.1.1 PBDE concentrations in sediment can be compared to OSPAR BACs and EAC which were primarily developed for the assessment of contaminant concentrations. BACs are used to make precautionary tests of whether observed concentrations are 'near background' or 'close to zero' in the case of man-made substances (i.e., PBDEs). PBDEs with concentrations below the BAC fulfil the ultimate aim of the OSPAR Hazardous Substances Strategy of approaching the natural Background Concentration (BC). EACs were developed to assess whether concentrations of contaminants are likely to have adverse biological effects on the marine environment. For PBDEs, the Canadian FEQGs adapted for use in CEMP assessments (OSPAR, 2020<sup>8</sup>) were used as EAC equivalents. Hence, adverse effects on marine organisms are rarely observed when concentrations are below the FEQG.

### 7.2 Caledonia North Site

7.2.1.1 Concentrations of PBDEs in sediment collected across the Caledonia North Site were recorded below the LOD with the exception of PBDE 209 which ranged from below the LOD to 0.37 ng/g (Table 7-1). The concentration of PBDE 209 at Station ENV35 was below the BAC (OSPAR, 2020<sup>8</sup>) and FEQG (Vinas *et al.*, 2023<sup>9</sup>) guidelines. Therefore, toxic effects to fauna by PBDEs in the Caledonia North Site are unlikely, even with the disturbance of sediment.

Table 7-1: Concentration of PBDE 209 identified when normalised to 2.5% TOC from sediment samples across the Caledonia North Site.

Sample Station	PBDE 209 (ng/g Dry Weight)
ENV07	<LOD
ENV18	<LOD
ENV33	<LOD
ENV35	0.37
BAC Value (OSPAR, 2020 <sup>8</sup> )	0.5
FEQG Value (Vinas <i>et al.</i> , 2023 <sup>9</sup> )	47.5

## 7.3 Caledonia North OECC

7.3.1.1 Contamination of PBDEs in sediment collected across the Caledonia North OECC were recorded below the LOD with the exception of PBDE 209 which ranged from below the LOD at Station ENV58 to 1.86ng/g at Station ENV31 (Table 7-2). Analysis of survey results showed that PBDE 209 at all but Station ENV58 exceeded the OSPAR (2020<sup>8</sup>) BAC. However, concentrations were all well below the FEQGs. Therefore, toxic effects to fauna by PBDEs in the Caledonia North OECC are unlikely, even with the disturbance of sediment.

Table 7-2: Concentration of PBDE 209 identified when normalised to 2.5% TOC from sediment samples across the Caledonia North OECC.

Sample Station	PBDE 209 (ng/g Dry Weight)
ENV04	0.80
ENV13	0.71
ENV17	0.61
ENV26	1.53
ENV29	0.28
ENV31	1.86
ENV36	0.39
ENV53	0.23
ENV58	<LOD
BAC Value (OSPAR, 2020 <sup>8</sup> )	0.5
FEQG Value (Vinas <i>et al.</i> , 2023 <sup>9</sup> )	47.5

## 8 Organochloropesticides

### 8.1 Overview

8.1.1.1 OCPs are synthetic pesticides widely used all over the world. They belong to the group of chlorinated hydrocarbon derivatives, which have vast application in the chemical industry and in agriculture. These compounds are known for their high toxicity, slow degradation, and bioaccumulation (Jayaraj *et al.*, 2017<sup>27</sup>).

### 8.2 Caledonia North Site

8.2.1.1 Results from a site-specific survey showed OCPs present in sediment samples collected across the Caledonia North Site were generally below the LOD, with the exception of p,p'-Dichlorodiphenyldichloroethane at Station ENV33 (0.1ng/g). There are currently no ALs for OCPs (Marine Scotland, 2017<sup>1</sup>) and toxic effects to fauna in the Caledonia North Site are not anticipated.

### 8.3 Caledonia North OECC

8.3.1.1 Results from a site-specific survey showed OCPs present in sediment samples collected across the Caledonia North OECC were generally below the LOD, with the exception of dieldrin at Station ENV04. There are currently no ALs for OCPs (Marine Scotland, 2017<sup>1</sup>) and toxic effects to fauna in the Caledonia North OECC are not anticipated.

## 9 Organotins

### 9.1 Overview

9.1.1.1 Organotin compounds are man-made metallic tin complexes with hydrocarbon substituents with historical applications in food packages, pesticides, wood preservatives, antifouling, and anticorrosion paints. According to the OSPAR Intermediate assessment, mean concentrations of TBT have measurably reduced in the Southern North Sea and are very low or undetectable elsewhere (OSPAR, 2017<sup>11</sup>).

### 9.2 Caledonia North Site

9.2.1.1 Organotin concentrations found in sediment samples collected across the Caledonia North Site were all below the LOD at all sampled stations. Therefore, toxic effects to fauna are not anticipated.

### 9.3 Caledonia North OECC

9.3.1.1 Organotin concentrations found in sediment samples across the Caledonia North OECC were all below the LOD at all sampled stations. Therefore, toxic effects to fauna are not anticipated.

## 10 Metals

### 10.1 Overview

10.1.1.1 Metals constitute a significant source of pollution in marine environments, originating from both natural and anthropogenic sources. Within the water column, biogeochemical processes govern the existence of heavy metals in particulate or dissolved form. Additionally, some proportion of metals get deposited in marine sediment which act as a major sink for metal pollution. Metals are taken up by marine biota through seawater or through food chain, which may adversely affect them by altering their abundance, diversity, and physiology. As detailed in paragraph 3.1.1.4 and subsequent sections, the EIAs for the Moray East, Moray West, and Beatrice OWFs identified no metals exceeding ALs, nor were any present in concentrations that would be likely to affect surrounding faunal communities.

### 10.2 Caledonia North Site

10.2.1.1 The results of the site-specific survey of the Caledonia North Site showed that chromium, copper, nickel and zinc were positively correlated with fines and negatively correlated with sands, meaning that fluctuations in metal concentrations were influenced by variations in sediment particle size and the resultant adsorption properties. These four metals additionally demonstrated positive correlations with TOC and N-alkalines. Alternatively, Mercury (Hg) was below the LOD in all samples. Concentrations of metals were all well below their ALs (Table 10-1), indicating that toxic impacts are unlikely to occur.

Table 10-1: Summary statistics of sediment metal contamination from samples collected across the Caledonia North Site.

Metal	AL1	AL2	Minimum	Maximum	Mean	Standard Deviation
Arsenic	20	70	3.7	6.9	5.3	1.7
Cadmium	0.4	4	0.09	0.12	0.11	0.001
Chromium	40	370	6	10.2	8	1.8
Copper	40	300	3	4.4	3.45	0.7
Mercury	0.3	1.5	NQ	NQ	NC	NC
Nickel	20	150	2.6	3.8	3.1	0.5
Lead	50	400	3.1	3.5	3.4	0.2
Zinc	130	600	6.3	13.1	8.15	3.3

Concentrations expressed as ug/g dry weight sediment. NQ – not qualified due to concentrations below the LOD. NC – Not calculated due to concentrations below the LOD.

### 10.3 Caledonia North OECC

10.3.1.1 The results of the site-specific survey of the Caledonia North OECC showed arsenic, cadmium, chromium, copper, mercury, nickel, lead and zinc were positively correlated with depth and finer sediment, indicating that fluctuations in metal concentrations were influenced by variations in sediment particle size and resultant adsorption properties. These five metals also demonstrated positive correlations with TOC and low molecular weight n-alkanes. Concentrations of metals were all well below their ALs (Table 10-2), indicating that toxic impacts are unlikely to occur.

Table 10-2: Summary statistics of sediment metal contamination from samples collected across the Caledonia North OECC.

Metal	AL1	AL2	Minimum	Maximum	Mean	Standard Deviation
Arsenic	20	70	2.1	5.4	3.2	1.1
Cadmium	0.4	4	NQ	0.12	NC	NC
Chromium	40	370	6.1	14.7	10.1	2.9
Copper	40	300	3.0	5.9	3.9	1.0
Mercury	0.3	1.5	NQ	0.01	NC	NC
Nickel	20	150	3.3	9.8	6.1	2.3
Lead	50	400	2.1	6.2	3.9	1.4
Zinc	130	600	10.6	24.8	15.8	4.6
Concentrations expressed as ug/g dry weight sediment. NQ – not qualified due to concentrations below the LOD. NC – Not calculated due to concentrations below the LOD.						

# 11 Sediment Contamination Conclusions

11.1.1.1 All contaminants were below their respective guideline thresholds (Table 11-1), frequently falling below the LOD. Consistent with findings from the Moray West, Moray East and Beatrice OWFs, the survey results indicate no evidence of unacceptable or significant contamination levels within the MW&SQ Study Area.

Table 11-1 Summary of sediment contamination in the Caledonia North Site and Caledonia North OECC relative to threshold guidelines.

Contaminant Type	Contamination Summary Relative to Threshold Guidelines
THC	No exceedance of AL1 (Marine Scotland, 2017 <sup>1</sup> ). No exceedance of threshold values (Kjeilen <i>et al.</i> , 2004 <sup>18</sup> ; UKOOA, 2002 <sup>19</sup> ; 2005 <sup>20</sup> ).
PAH	No exceedance of: <ul style="list-style-type: none"> <li>▪ TEL/PEL values (CCME, 1999<sup>5</sup>)</li> <li>▪ ERL/ERM values (Long <i>et al.</i>, 1995<sup>25</sup>)</li> </ul>
PCB	No exceedance of AL1/AL2 (Marine Scotland, 2017 <sup>1</sup> ). Below the LOD at all stations.
PBDE	All PBDEs were recorded below the LOD, except PBDE 209, which was: <ul style="list-style-type: none"> <li>▪ Below the BAC (OSPAR, 2020a<sup>7</sup>) and FEQGs (OSPAR, 2020b<sup>8</sup>) across the Caledonia OWF</li> <li>▪ Above the BAC at four Stations in the Caledonia North OECC but below FEQGs.</li> </ul>
OCP	There are currently no ALs for OCPs (Marine Scotland, 2017 <sup>1</sup> ). Below the LOD at most stations.
Organotins	No exceedance of AL1/AL2 (Marine Scotland, 2017 <sup>1</sup> ). Below the LOD at all stations.
Metals	No exceedance of AL1/AL2 (Marine Scotland, 2017 <sup>1</sup> ).

## 12 References

<sup>1</sup>Marine Scotland (2017) 'Pre-disposal sampling guidance'. Available at:

<https://www.gov.scot/binaries/content/documents/govscot/publications/advice-and-guidance/2020/02/marine-licensing-applications-and-guidance/documents/guidance/pre-disposal-sampling-guidance/pre-disposal-sampling-guidance/govscot%3Adocument/Pre-disposal%2Bsampling%2Bguidance.pdf> (Accessed 20/05/2024)

<sup>2</sup> International Maritime Organization (IMO) (2014) 'Waste Assessment Guidelines under the London Convention and Protocol'. IMO: London, UK. ISBN 978-92-801-1613-7

<sup>3</sup> OSPAR Commission (2014) 'Guidelines for the Management of Dredged Material at Sea'. Available at: <https://www.ospar.org/documents?d=34060> (Accessed 20/05/2024)

<sup>4</sup> Gorham-Test, C., Wade, T., Engle, V., Summers, K. and Hornig, E. (1999) 'Regional Environmental Monitoring and Assessment Program — Galveston Bay 1993'. Proceedings, Galveston Bay Estuary Program, State of the Bay Symposium IV, January 28–29, Galveston, TX, 97–109

<sup>5</sup> Canadian Council of Ministers of the Environment (1999) 'Canadian sediment quality guidelines for the protection of aquatic life: Polycyclic aromatic hydrocarbons (PAHs)'. Available at: <https://ccme.ca/en/res/polycyclic-aromatic-hydrocarbons-pahs-canadian-sediment-quality-guidelines-for-the-protection-of-aquatic-life-en.pdf> (Accessed 20/05/2024)

<sup>6</sup> OSPAR Commission (2023) 'Levels and trends in marine contaminants and their biological effects – CEMP Assessment report 2023'. Available at: <https://oap.ospar.org/en/ospar-assessments/committee-assessments/hazardous-substances-and-eutrophication/mime/cemp-levels-and-trends-marine-contaminants/cemp-2023/> (Accessed 21/05/2024)

<sup>7</sup> OSPAR Commission (2020a) 'Background document for Canadian Federal Environmental Quality Guidelines (FEQGs) for Polybrominated Diphenyl Ethers (PBDEs) in sediment and biota', Hazardous substances and Eutrophication Series. Available at: <https://www.ospar.org/documents?v=42746> (Accessed 20/05/2024)

<sup>8</sup> OSPAR Commission (2020b) 'Background document for Canadian Federal Environmental Quality Guidelines (FEQGs) for Polybrominated Diphenyl Ethers (PBDEs) in sediment and biota', Hazardous substances and Eutrophication Series. Available at: <https://www.ospar.org/documents?v=42746> (Accessed 20/05/2024)

<sup>9</sup> Viñas, L., Soerensen, A.L. and Fryer, R. (2023) 'Status and trends of polybrominated diphenyl ethers (PBDEs) in Biota and sediment, OSPAR'. Available at: <https://oap.ospar.org/en/ospar-assessments/quality-status-reports/qsr-2023/indicator->

[assessments/status-and-trends-polybrominated-diphenyl-ethers-pbdes-biota-and/](#)  
(Accessed 20/05/2024)

<sup>10</sup> Davies, J., Baxter, J., Bradley, M., Connor, D., Khan, J., Murray, E., Sanderson, W., Turnbull, C. and Vincent, M. (eds). (2001). Marine Monitoring Handbook, JNCC, Peterborough

<sup>11</sup> OSPAR (2017) 'Intermediate Assessment 2017'. Available at:  
<https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/> (Accessed 30/05/2024)

<sup>12</sup> Moray Offshore Windfarm (West) Limited. (2018). Moray West Offshore Wind Farm Environmental Statement. Available at: <https://www.moraywest.com/documentlibrary> (Accessed 21/05/2024)

<sup>13</sup> Moray Offshore Renewables Limited (2012a) 'Moray East OWF Environmental Statement (Volume 10, Appendix 4.2 A: Benthic Ecology Characterisation Survey (Wind Farm Sites))'. Available at: <https://www.morayeast.com/application/files/1315/8014/0645/Appendix-4-2-A-Benthic-Ecology-Wind-Farm-Sites.pdf> (Accessed 21/05/24)

<sup>14</sup> Moray Offshore Renewables Limited (2012b) 'Moray East OWF Environmental Statement (Volume 2, Chapter 4: Biological Environment Baseline)'. Available at:  
[https://marine.gov.scot/sites/default/files/chapter\\_4\\_-\\_biological\\_environment\\_baseline.pdf](https://marine.gov.scot/sites/default/files/chapter_4_-_biological_environment_baseline.pdf) (Accessed 30/05/2024)

<sup>15</sup> Beatrice Offshore Wind Farm Limited (BOWL). (2012) 'Beatrice OWF Environmental Statement'. Available at: <https://marine.gov.scot/data/environmental-statement-construction-operation-generating-station-and-transmission-works> (Accessed 21/05/2024)

<sup>16</sup> Repsol Sinopec Resources UK Limited (2018) 'Beatrice Field Decommissioning Environmental Impact Assessment'. Available at:  
[https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/731309/Beatrice\\_Environmental\\_Assessment\\_Report.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/731309/Beatrice_Environmental_Assessment_Report.pdf) (Accessed 21/05/2024)

<sup>17</sup> McDougall, J. (2000) 'Section 5.1. The significance of hydrocarbons in surficial sediments from the Atlantic Margin regions'. In Hydrocarbons in environmental surveys of the seafloor of the UK Atlantic Margin. Daventry, Northants NN11 5EA, UK.: Geotek Limited ISBN 09538399-0-7

<sup>18</sup> Kjeilen-Eilertsen, G., Westerlund, S., Bamber, S., Tandber, A.H., Myhre, L.P. and Tvedten, O. (2004) 'UKOOA phase III- Characterisation of Beryl, Brent A, Brent S, Clyde and Miller cuttings piles through field work, laboratory studies and chemical analysis'. Final Report – 2004-197

<sup>19</sup> UKOOA (2002) 'UKOOA Drill Cuttings Initiative Final Report'. February 2002. UKOOA Drill Cuttings Initiative Executive Committee

- <sup>20</sup> UKOOA (2005) 'UKOOAJIP 2004 Drill Cuttings Initiative Phase III'. Final Report. 20132900
- <sup>21</sup> Kingston, P.F. (1992) 'Impact of offshore oil production installations on the benthos of the North Sea'. ICES Journal of Marine Science 49(1): 45-53. Available at: <https://academic.oup.com/icesjms/article/49/1/45/732779> (Accessed 21/05/2024)
- <sup>22</sup> Bouloubassi, I., Fillaux, J. and Saliot, A. (2001) 'Hydrocarbons in surface sediments from the Changjiang (Yangtze River) Estuary, East China Sea'. Marine Pollution Bulletin, : 1335-46.
- <sup>23</sup> Harborne, J.B. (1999) 'Phytochemical Dictionary: A Handbook of Bioactive Compounds from Plants'. Boca Raton, Florida, USA: CRC Press
- <sup>24</sup> Tran, K., Yu, C. and Zeng, E. (1995) 'Petrogenic and Biogenic Sources of N-alkanes off San Diego, California'. Available at: [ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/1994\\_95AnnualReport/ar05.pdf](ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/1994_95AnnualReport/ar05.pdf) (Accessed 21/05/2024)
- <sup>25</sup> Long, E.R., MacDonald, D.D., Smith, S.L. and Calder, F.D. (1995) 'Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments'. Environmental Management 19: 81-97
- <sup>26</sup> Marine Scotland (2020) 'Contaminant and biological effect data 1999-2018 for the National Performance Framework Clean Seas Indicator 2019 and Scotland's Marine Assessment 2020'
- <sup>27</sup> Jayaraj, R., Megha, P. and Sreedev, P. (2016) 'Review article. Organochlorine pesticides, their toxic effects on living organisms and their fate in the environment', Interdisciplinary Toxicology 9(3-4): 90-100

Caledonia Offshore Wind Farm  
5th Floor, Atria One  
144 Morrison Street  
Edinburgh  
EH3 8EX

[www.caledoniaoffshorewind.com](http://www.caledoniaoffshorewind.com)

