



Annex 11.2G: Density estimates per survey of birds in flight

MachairWind Offshore Ornithology

ScottishPower Renewables (SPR)

320 St Vincent St Glasgow G2 5AD

Prepared by:

SLR Consulting Limited

St. Vincent Place, Glasgow, G1 2EU

SLR Project No.: 413.058018.00001

Client Reference No: MCW-DWF-ENV-REP-RHS-000199

2 April 2026

Revision: 3

Revision Record

Revision	Date	Prepared By	Checked By	Authorised By
Rev 1	17 November 2025	NG & DJ	NG/LS	ES
Rev 2	26 February 2026	DJ	NG/LS	ES
Rev 3	2 April 2026	DJ	NG/LS	ES



Table of Contents

1.0 Introduction	1
2.0 Density estimates.....	2
2.1 Arctic Skua	4
2.2 Arctic Tern.....	6
2.3 Auk species.....	10
2.4 Auk/Shearwater species	12
2.5 Black-headed Gull.....	14
2.6 Black Guillemot	18
2.7 Common Tern.....	20
2.8 Common Gull	22
2.9 Common Tern	26
2.10 Cormorant/Shag.....	30
2.11 Diver species.....	32
2.12 Dunlin	34
2.13 European Storm-petrel.....	36
2.14 Fulmar	38
2.15 Fulmar/Gull species	40
2.16 Gannet	42
2.17 Golden Plover	44
2.18 Great Black-backed Gull	46
2.19 Great Northern Diver.....	50
2.20 Great Shearwater.....	52
2.21 Great Skua	54
2.22 Grey Phalarope	56
2.23 Guillemot	58
2.24 Guillemot/Razorbill	64
2.25 Gull species.....	66
2.26 Herring Gull	68
2.27 Kittiwake.....	72
2.28 Large Gull species	76
2.29 Lesser Black-backed Gull	78
2.30 Little Egret.....	82
2.31 Manx Shearwater	84
2.32 Puffin	88
2.33 Razorbill	94



2.34 Red-throated Diver.....	100
2.35 Small Gull species.....	102
2.36 Sooty Shearwater.....	104
2.37 Unidentified Bird species.....	106
2.38 Wader species	108

Tables in Text

Table 1: List of 30 digital aerial survey (DAS) samples recorded by the Project or Third-Party source. The survey date, survey number and a sample identification (ID) number is provided for each DAS sample.....	2
Table 2: Arctic Skua - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	4
Table 3: Arctic Tern - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	6
Table 4: ArcticTernApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	8
Table 5: Auk species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	10
Table 6: Auk/Shearwater species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	12
Table 7: Black-headed Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	14
Table 8: BlackHeadedGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	16



Table 9:	Black Guillemot - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	18
Table 10:	Commic Tern - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	20
Table 11:	Common Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	22
Table 12:	CommonGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	24
Table 13:	Common Tern - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	26
Table 14:	CommonTernApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	28
Table 15:	Cormorant/Shag - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	30
Table 16:	Diver species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	32
Table 17:	Dunlin - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	34
Table 18:	European Storm-petrel Apportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	36



Table 19: Fulmar - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	38
Table 20: Fulmar/Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	40
Table 21: Gannet - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	42
Table 22: Golden Plover - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	44
Table 23: Great Black-backed Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	46
Table 24: GreatBlackBackedGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	48
Table 25: Great Northern Diver - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	50
Table 26: Great Shearwater - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	52
Table 27: Great Skua - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	54
Table 28: Grey Phalarope - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	56



Table 29: Guillemot - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	58
Table 30: GuillemotApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	60
Table 31: GuillemotApportionedCorrected - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	62
Table 32: Guillemot/Razorbill - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	64
Table 33: Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	66
Table 34: Herring Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	68
Table 35: HerringGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	70
Table 36: Kittiwake - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	72
Table 37: KittiwakeApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	74
Table 38: Large Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	76



Table 39: Lesser Black-backed Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	78
Table 40: LesserBlackBackedGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	80
Table 41: Little Egret - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	82
Table 42: Manx Shearwater - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	84
Table 43: ManxShearwaterApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	86
Table 44: Puffin - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	88
Table 45: PuffinApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	90
Table 46: PuffinApportionedCorrected - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	92
Table 47: Razorbill - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	94
Table 48: RazorbillApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	96



Table 49: RazorbillApportionedCorrected - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	98
Table 50: Red-throated Diver - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	100
Table 51: Small Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	102
Table 52: Sooty Shearwater - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	104
Table 53: Unidentified Bird species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).....	106
Table 54: Wader species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).	108



1.0 Introduction

1. This **Annex 11.2G: Density estimates per survey of birds in flight** provides density estimates of all birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer used to inform the baseline site characterisation for the Project.
2. Density estimates were calculated for 30 Digital Aerial Survey (DAS) samples (one sample is composed of one complete DAS survey per month) which are summarised in **Table 1**. The DAS samples were selected from Project DAS surveys and Third-Party DAS surveys; refer to **Technical Appendix 11.2: Baseline Site Characterisation** for an explanation of survey areas. It was agreed with NatureScot to use the 30 selected DAS samples presented to inform the baseline site characterisation for the Project (NatureScot letter dated 7 August 2025).
3. This **Annex 11.2G** provides density estimates of all birds in flight identified to species and birds assigned to a 'species group' as well as for the total number of birds calculated after species groups had been assigned to a specific species (for an explanation of apportioning birds to a species from a broader species group, refer to **Technical Appendix 11.2: Baseline Site Characterisation**). In addition, for guillemots, razorbills and puffins, density estimates include birds which were 'corrected' because they were under the water at the time of each DAS; 'corrected' guillemots, razorbills and puffins are added together with apportioned guillemots, razorbills and puffins to provide a total density estimate for each of these species (for an explanation of how guillemots, razorbills and puffins were 'corrected', refer to **Technical Appendix 11.2: Baseline Site Characterisation**). Note, tables summarising both apportioned and apportioned plus dive bias corrected numbers are presented. Dive bias corrected numbers are only applied to counts of birds on the sea surface, and thus for guillemots, razorbills and puffins, birds in flight in the 'apportioned and dive bias corrected' tables have the same numbers as birds in flights in the 'apportioned only' tables.
4. In the table titles introducing **Table 2** to **Table 54**, birds with 'Apportioned' affixed to the species name are species which include apportioned birds. For example, 'Guillemot' refers to density estimates based solely on positively identified guillemots, whereas 'GuillemotApportioned' refers to density estimates derived from all positively identified guillemots plus individuals from apportioned species groups such as 'Auk Species' and 'Guillemot/Razorbill'. Similarly, 'Corrected' in table titles means that guillemot, razorbill and puffin density estimates are adjusted to account for birds under the water. For guillemot, razorbill and puffin, the 'ApportionedCorrected' tables provide the total density estimates for these species.



2.0 Density estimates

5. Density estimates for all species in the WDA, WDA plus 2 km buffer and WDA plus 4 km buffer in 30 DAS samples are presented in **Table 2** to **Table 54**. For an explanation of how density was calculated, refer to **Technical Appendix 11.2: Baseline Site Characterisation**.

Table 1: List of 30 digital aerial survey (DAS) samples recorded by the Project or Third-Party source. The survey date, survey number and a sample identification (ID) number is provided for each DAS sample.

Date of DAS survey	Project or Third-Party DAS (survey number)	ID sample number
21st March 2021	Third-Party (6)	1
20th April 2021	Project (1)	2
11th May 2021	Project (2)	3
12th June 2021	Project (3)	4
6th July 2021	Project (4)	5
14th August 2021	Project (5)	6
29th September 2021	Project (6)	7
22nd October 2021	Project (7)	8
23rd November 2021	Project (8)	9
15th December 2021	Third-Party (15)	10
20th January 2022	Project (9)	11
1st plus 4th March 2021 (proxy sample for February 2022)	Third-Party (5)	12
28th March 2022	Project (11)	13
24th April 2022	Project (12)	14
May 28th 2022	Project (13)	15
June 2022	No sample	N/A
27th July 2022	Project (15)	16
21st August 2022	Project (16)	17
13th September 2022	Project (17)	18
4th November 2022 (proxy sample for October 2022)	Project (18)	19
22nd November 2022	Project (19)	20
2nd December 2022	Project (20)	21
28th January 2023	Project (22)	22
24th February 2023	Project (23)	23
19th plus 23rd March 2023	Project (24)	24
22nd April 2023	Project (25)	25
13th May 2023	Project (26)	26
07th June 2023	Project (27)	27



Date of DAS survey	Project or Third-Party DAS (survey number)	ID sample number
21st July 2023	Project (28)	28
16th August 2023	Project (29)	29
06th September 2023	Project (30)	30



2.1 Arctic Skua

Table 2: Arctic Skua - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.03	0.03 (0.03)	0.00–0.08	1	0.02	0.02 (0.02)	0.00–0.06	1	0.01	0.01 (0.02)	0.00–0.06	2
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.2 Arctic Tern

Table 3: Arctic Tern - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.02	0.02 (0.02)	0.00–0.07	1.00
3	0.11	0.11 (0.04)	0.04–0.18	0.36	0.09	0.08 (0.03)	0.03–0.14	0.38	0.07	0.08 (0.03)	0.03–0.13	0.38
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.02	0.02 (0.02)	0.00–0.05	1.00
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
16	0.08	0.08 (0.05)	0.00–0.18	0.62	0.08	0.08 (0.03)	0.02–0.15	0.38	0.07	0.08 (0.03)	0.03–0.13	0.38
17	0.14	0.14 (0.06)	0.03–0.27	0.43	0.20	0.19 (0.07)	0.07–0.34	0.37	0.27	0.27 (0.10)	0.10–0.50	0.37



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.06	0.06 (0.04)	0.00–0.16	0.67	0.05	0.05 (0.04)	0.00–0.13	0.80
27	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
28	0.03	0.03 (0.02)	0.00–0.09	0.67	0.05	0.05 (0.03)	0.00–0.11	0.60	0.04	0.04 (0.02)	0.00–0.09	0.50
29	0.07	0.07 (0.05)	0.00–0.19	0.71	0.05	0.05 (0.03)	0.00–0.13	0.60	0.08	0.08 (0.04)	0.02–0.16	0.50
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Table 4: ArcticTernApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.02	0.02 (0.02)	0.00–0.07	1.00
3	0.25	0.25 (0.07)	0.12–0.39	0.28	0.19	0.19 (0.05)	0.09–0.29	0.26	0.16	0.17 (0.04)	0.09–0.25	0.24
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.02	0.02 (0.02)	0.00–0.05	1.00
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.01	0.01 (0.01)	0.00–0.02	1.00	0.02	0.02 (0.01)	0.01–0.05	0.50	0.02	0.02 (0.01)	0.01–0.04	0.50
7	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.01	0.01 (0.01)	0.00–0.04	1.00
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.01	0.01 (0.01)	0.00–0.02	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
15	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00
16	0.15	0.16 (0.07)	0.03–0.32	0.44	0.13	0.13 (0.05)	0.05–0.23	0.38	0.12	0.12 (0.04)	0.04–0.20	0.33
17	0.27	0.27 (0.08)	0.12–0.45	0.30	0.38	0.38 (0.09)	0.23–0.57	0.24	0.42	0.42 (0.11)	0.24–0.65	0.26



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.06	0.06 (0.04)	0.00–0.16	0.67	0.05	0.05 (0.04)	0.00–0.13	0.80
27	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
28	0.03	0.03 (0.02)	0.00–0.09	0.67	0.05	0.05 (0.03)	0.00–0.11	0.60	0.04	0.04 (0.02)	0.00–0.09	0.50
29	0.07	0.07 (0.05)	0.00–0.19	0.71	0.05	0.05 (0.03)	0.00–0.13	0.60	0.08	0.08 (0.04)	0.02–0.16	0.50
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.3 Auk species

Table 5: Auk species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.06	0.06 (0.03)	0.00–0.13	0.5	0.04	0.04 (0.03)	0.00–0.09	0.75	0.04	0.04 (0.02)	0.01–0.09	0.5
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.0
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.0
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.4 Auk/Shearwater species

Table 6: Auk/Shearwater species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.01	0.01 (0.01)	0.00–0.04	1.00	0.02	0.02 (0.01)	0.00–0.05	0.50	0.01	0.01 (0.01)	0.00–0.04	1.00
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
6	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.03	0.03 (0.02)	0.01–0.07	0.67
7	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.01	0.01 (0.02)	0.00–0.04	2.00	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.01 (0.01)	0.00–0.04	1.00
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.04	0.04 (0.02)	0.00–0.10	0.50	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.04	0.50
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.04	0.04 (0.03)	0.00–0.11	0.75	0.04	0.04 (0.02)	0.00–0.10	0.50	0.03	0.03 (0.02)	0.00–0.07	0.67
25	0.07	0.07 (0.04)	0.01–0.16	0.57	0.05	0.05 (0.03)	0.01–0.11	0.60	0.08	0.08 (0.04)	0.02–0.15	0.50
26	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
29	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.02 (0.01)	0.01–0.04	0.50
30	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.04	0.50



2.5 Black-headed Gull

Table 7: Black-headed Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Table 8: BlackHeadedGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



2.6 Black Guillemot

Table 9: Black Guillemot - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



2.7 Commic Tern

Table 10: Commic Tern - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.14	0.14 (0.06)	0.04–0.26	0.43	0.10	0.10 (0.04)	0.03–0.19	0.40	0.09	0.09 (0.03)	0.04–0.16	0.33
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.13	0.12 (0.05)	0.04–0.22	0.42	0.13	0.13 (0.05)	0.06–0.23	0.38	0.11	0.11 (0.04)	0.05–0.19	0.36
7	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.01	0.01 (0.01)	0.00–0.04	1.00
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
15	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
16	0.08	0.08 (0.06)	0.00–0.22	0.75	0.07	0.07 (0.04)	0.01–0.16	0.57	0.05	0.05 (0.03)	0.00–0.13	0.60
17	0.14	0.14 (0.05)	0.04–0.25	0.36	0.19	0.19 (0.06)	0.09–0.30	0.32	0.16	0.15 (0.05)	0.07–0.24	0.33



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.8 Common Gull

Table 11: Common Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.01	0.01 (0.01)	0.00–0.04	1.00	0.05	0.05 (0.03)	0.00–0.13	0.60	0.05	0.05 (0.03)	0.01–0.11	0.6
10	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.04	0.5
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
20	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.04	0.05 (0.02)	0.01–0.10	0.4
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
23	0.03	0.03 (0.02)	0.00–0.07	0.67	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.5
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.0
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Table 12: CommonGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.01	0.01 (0.01)	0.00–0.04	1.00	0.05	0.05 (0.03)	0.00–0.13	0.60	0.05	0.05 (0.03)	0.01–0.11	0.6
10	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.04	0.5
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
20	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.04	0.05 (0.02)	0.01–0.10	0.4
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
23	0.03	0.03 (0.02)	0.00–0.07	0.67	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.5
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.0
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.9 Common Tern

Table 13: Common Tern - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.0
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.11	0.11 (0.11)	0.00–0.34	1	0.08	0.08 (0.08)	0.00–0.23	1.0	0.06	0.06 (0.06)	0.00–0.18	1.0
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1.0	0.01	0.01 (0.01)	0.00–0.02	1.0
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.01	0.01 (0.01)	0.00–0.04	1	0.02	0.02 (0.01)	0.00–0.05	0.5	0.01	0.02 (0.01)	0.00–0.04	0.5



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Table 14: CommonTernApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.23	0.22 (0.12)	0.05–0.49	0.55	0.20	0.20 (0.09)	0.06–0.39	0.45	0.16	0.16 (0.07)	0.05–0.32	0.44
7	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.01	0.01 (0.01)	0.00–0.04	1.00
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.02	0.02 (0.02)	0.00–0.06	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.03	0.03 (0.02)	0.00–0.06	0.67	0.03	0.03 (0.01)	0.00–0.06	0.33	0.02	0.02 (0.01)	0.00–0.05	0.50
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.10 Cormorant/Shag

Table 15: Cormorant/Shag - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.11 Diver species

Table 16: Diver species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



2.12 Dunlin

Table 17: Dunlin - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



2.13 European Storm-petrel

Table 18: European Storm-petrel Apportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.05	0.05 (0.04)	0.00–0.14	0.80
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.08	0.08 (0.03)	0.03–0.15	0.38	0.13	0.13 (0.04)	0.07–0.21	0.31	0.12	0.12 (0.03)	0.06–0.19	0.25



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.04	0.04 (0.04)	0.00–0.11	1.00	0.03	0.03 (0.03)	0.00–0.09	1.00
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.01	0.01 (0.01)	0.00–0.04	1.00	0.03	0.03 (0.02)	0.00–0.06	0.67	0.03	0.03 (0.02)	0.00–0.07	0.67
28	0.23	0.23 (0.07)	0.10–0.36	0.30	0.30	0.30 (0.06)	0.17–0.44	0.20	0.27	0.27 (0.06)	0.16–0.38	0.22
29	0.00	0.00 (0.00)	0.00–0.00		0.03	0.03 (0.01)	0.00–0.06	0.33	0.15	0.15 (0.05)	0.06–0.25	0.33
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.14 Fulmar

Table 19: Fulmar - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.04	0.04 (0.02)	0.01–0.07	0.50	0.05	0.05 (0.02)	0.02–0.08	0.40	0.07	0.07 (0.01)	0.04–0.10	0.14
2	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
3	0.07	0.07 (0.03)	0.01–0.14	0.43	0.06	0.06 (0.02)	0.02–0.10	0.33	0.05	0.05 (0.02)	0.01–0.09	0.40
4	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
5	0.04	0.04 (0.02)	0.01–0.08	0.50	0.04	0.04 (0.02)	0.01–0.08	0.50	0.03	0.03 (0.01)	0.01–0.06	0.33
6	0.07	0.07 (0.04)	0.00–0.15	0.57	0.10	0.09 (0.03)	0.04–0.17	0.33	0.08	0.08 (0.03)	0.03–0.14	0.38
7	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
8	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
9	0.10	0.10 (0.03)	0.04–0.14	0.30	0.12	0.12 (0.03)	0.07–0.17	0.25	0.10	0.10 (0.02)	0.05–0.15	0.20
10	0.09	0.08 (0.02)	0.04–0.11	0.25	0.10	0.09 (0.02)	0.05–0.14	0.22	0.10	0.10 (0.02)	0.07–0.13	0.20
11	0.42	0.42 (0.23)	0.07–0.95	0.55	0.30	0.30 (0.17)	0.05–0.67	0.57	0.30	0.31 (0.15)	0.10–0.65	0.48
12	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.04	0.50
13	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.04	0.04 (0.01)	0.02–0.07	0.25
14	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.05	0.50
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.06	0.05 (0.03)	0.00–0.13	0.60	0.05	0.05 (0.02)	0.01–0.09	0.40	0.04	0.04 (0.02)	0.01–0.07	0.50



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.04	0.04 (0.02)	0.00–0.10	0.50	0.04	0.04 (0.02)	0.01–0.07	0.50	0.05	0.05 (0.02)	0.01–0.10	0.40
18	0.07	0.06 (0.03)	0.01–0.13	0.50	0.13	0.13 (0.08)	0.02–0.31	0.62	0.12	0.12 (0.06)	0.02–0.27	0.50
19	0.01	0.01 (0.01)	0.00–0.04	1.00	0.05	0.05 (0.02)	0.01–0.10	0.40	0.09	0.09 (0.03)	0.03–0.15	0.33
20	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
21	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.01	0.01 (0.01)	0.00–0.04	1.00
22	0.08	0.08 (0.04)	0.03–0.15	0.50	0.07	0.07 (0.03)	0.02–0.12	0.43	0.05	0.05 (0.02)	0.02–0.11	0.40
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.10	0.10 (0.03)	0.04–0.15	0.30	0.10	0.09 (0.03)	0.05–0.14	0.33	0.12	0.12 (0.03)	0.07–0.18	0.25
25	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.05	0.05 (0.02)	0.02–0.08	0.40
26	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.05	0.50
27	0.03	0.03 (0.02)	0.00–0.07	0.67	0.04	0.04 (0.02)	0.00–0.08	0.50	0.03	0.03 (0.01)	0.01–0.06	0.33
28	0.07	0.07 (0.03)	0.01–0.14	0.43	0.07	0.07 (0.03)	0.02–0.13	0.43	0.10	0.10 (0.04)	0.04–0.18	0.40
29	0.06	0.06 (0.03)	0.01–0.12	0.50	0.04	0.04 (0.02)	0.01–0.08	0.50	0.03	0.03 (0.02)	0.01–0.06	0.67
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.15 Fulmar/Gull species

Table 20: Fulmar/Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.16 Gannet

Table 21: Gannet - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.02)	0.00–0.05	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
2	0.08	0.08 (0.04)	0.03–0.17	0.50	0.08	0.09 (0.03)	0.04–0.14	0.33	0.15	0.15 (0.04)	0.07–0.25	0.27
3	0.12	0.12 (0.05)	0.04–0.24	0.42	0.16	0.16 (0.04)	0.09–0.26	0.25	0.16	0.16 (0.04)	0.09–0.25	0.25
4	0.15	0.16 (0.09)	0.03–0.35	0.56	0.15	0.15 (0.06)	0.05–0.29	0.40	0.19	0.19 (0.06)	0.08–0.32	0.32
5	0.17	0.16 (0.05)	0.07–0.26	0.31	0.18	0.18 (0.05)	0.09–0.28	0.28	0.15	0.15 (0.04)	0.08–0.23	0.27
6	0.20	0.15 (0.05)	0.06–0.27	0.33	0.14	0.14 (0.04)	0.07–0.23	0.29	0.14	0.14 (0.04)	0.08–0.22	0.29
7	0.23	0.23 (0.06)	0.11–0.36	0.26	0.18	0.18 (0.05)	0.09–0.28	0.28	0.14	0.14 (0.04)	0.07–0.22	0.29
8	0.07	0.07 (0.03)	0.01–0.14	0.43	0.06	0.06 (0.03)	0.02–0.11	0.50	0.09	0.09 (0.04)	0.02–0.19	0.44
9	0.03	0.03 (0.02)	0.00–0.07	0.67	0.04	0.04 (0.02)	0.01–0.08	0.50	0.06	0.06 (0.02)	0.02–0.11	0.33
10	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00
13	0.10	0.10 (0.04)	0.03–0.19	0.40	0.10	0.10 (0.04)	0.04–0.17	0.40	0.11	0.11 (0.03)	0.05–0.17	0.27
14	0.14	0.14 (0.05)	0.06–0.23	0.36	0.12	0.12 (0.04)	0.06–0.20	0.33	0.11	0.11 (0.03)	0.06–0.17	0.27
15	0.59	0.59 (0.17)	0.32–0.95	0.29	0.61	0.61 (0.13)	0.38–0.88	0.21	0.53	0.53 (0.10)	0.36–0.75	0.19
16	0.03	0.03 (0.03)	0.00–0.08	1.00	0.05	0.05 (0.03)	0.00–0.10	0.60	0.04	0.04 (0.02)	0.01–0.09	0.50



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.10	0.10 (0.04)	0.03–0.19	0.40	0.09	0.09 (0.03)	0.04–0.17	0.33	0.08	0.08 (0.03)	0.04–0.13	0.38
18	0.11	0.11 (0.03)	0.04–0.19	0.27	0.18	0.18 (0.04)	0.11–0.25	0.22	0.16	0.16 (0.03)	0.10–0.23	0.19
19	0.08	0.07 (0.05)	0.00–0.18	0.71	0.08	0.08 (0.04)	0.01–0.16	0.50	0.06	0.06 (0.03)	0.01–0.13	0.50
20	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
21	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
22	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
23	0.03	0.03 (0.01)	0.00–0.05	0.33	0.05	0.05 (0.02)	0.02–0.07	0.40	0.05	0.05 (0.02)	0.02–0.08	0.40
24	0.07	0.07 (0.03)	0.01–0.13	0.43	0.06	0.06 (0.02)	0.02–0.11	0.33	0.05	0.05 (0.02)	0.02–0.08	0.40
25	0.06	0.06 (0.04)	0.00–0.14	0.67	0.09	0.09 (0.03)	0.03–0.16	0.33	0.07	0.07 (0.02)	0.02–0.12	0.29
26	0.11	0.11 (0.05)	0.03–0.23	0.45	0.11	0.11 (0.04)	0.04–0.19	0.36	0.08	0.09 (0.03)	0.03–0.15	0.33
27	0.23	0.23 (0.07)	0.11–0.37	0.30	0.21	0.21 (0.05)	0.12–0.32	0.24	0.28	0.28 (0.06)	0.17–0.41	0.21
28	0.06	0.06 (0.02)	0.01–0.10	0.33	0.08	0.08 (0.04)	0.02–0.16	0.50	0.07	0.07 (0.03)	0.02–0.14	0.43
29	0.24	0.24 (0.06)	0.14–0.36	0.25	0.19	0.19 (0.05)	0.10–0.28	0.26	0.16	0.16 (0.04)	0.09–0.23	0.25
30	0.33	0.32 (0.08)	0.19–0.51	0.25	0.30	0.30 (0.06)	0.19–0.44	0.20	0.28	0.29 (0.06)	0.18–0.41	0.21



2.17 Golden Plover

Table 22: Golden Plover - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.18 Great Black-backed Gull

Table 23: Great Black-backed Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.07	0.07 (0.03)	0.01–0.13	0.43	0.14	0.14 (0.05)	0.05–0.25	0.36	0.12	0.12 (0.04)	0.05–0.21	0.33
10	0.09	0.08 (0.03)	0.03–0.14	0.38	0.06	0.06 (0.03)	0.02–0.12	0.50	0.08	0.08 (0.03)	0.03–0.15	0.38
11	0.01	0.01 (0.01)	0.00–0.04	1.00	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.02)	0.00–0.06	1.00
12	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.03	0.03 (0.01)	0.00–0.06	0.33
13	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.02 (0.01)	0.00–0.04	0.50
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
20	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00
21	0.03	0.03 (0.02)	0.00–0.07	0.67	0.03	0.03 (0.01)	0.00–0.06	0.33	0.03	0.03 (0.01)	0.01–0.05	0.33
22	0.04	0.04 (0.03)	0.00–0.11	0.75	0.03	0.03 (0.02)	0.00–0.08	0.67	0.02	0.02 (0.02)	0.00–0.06	1.00
23	0.04	0.04 (0.02)	0.01–0.08	0.50	0.05	0.05 (0.02)	0.02–0.08	0.40	0.04	0.04 (0.01)	0.01–0.07	0.25
24	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00



Table 24: GreatBlackBackedGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.08	0.08 (0.03)	0.03–0.14	0.38	0.14	0.14 (0.05)	0.05–0.26	0.36	0.13	0.13 (0.04)	0.05–0.22	0.31
10	0.09	0.09 (0.03)	0.03–0.15	0.33	0.07	0.07 (0.03)	0.02–0.12	0.43	0.09	0.09 (0.03)	0.03–0.15	0.33
11	0.01	0.01 (0.01)	0.00–0.04	1.00	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.02)	0.00–0.06	1.00
12	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.03	0.03 (0.01)	0.00–0.06	0.33
13	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.02 (0.01)	0.00–0.04	0.50
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
20	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00
21	0.03	0.03 (0.02)	0.00–0.07	0.67	0.03	0.03 (0.01)	0.00–0.06	0.33	0.03	0.03 (0.01)	0.01–0.05	0.33
22	0.04	0.04 (0.03)	0.00–0.11	0.75	0.03	0.03 (0.02)	0.00–0.08	0.67	0.02	0.02 (0.02)	0.00–0.06	1.00
23	0.04	0.04 (0.02)	0.01–0.08	0.50	0.05	0.05 (0.02)	0.02–0.08	0.40	0.04	0.04 (0.01)	0.01–0.07	0.25
24	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00



2.19 Great Northern Diver

Table 25: Great Northern Diver - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.20 Great Shearwater

Table 26: Great Shearwater - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
21	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.21 Great Skua

Table 27: Great Skua - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.02	0.02 (0.02)	0.00–0.06	1.00	0.02	0.02 (0.01)	0.00–0.05	0.50
5	0.01	0.01 (0.01)	0.00–0.04	1.0	0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.01 (0.01)	0.00–0.04	1.00
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.01	0.01 (0.01)	0.00–0.04	1.0	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.02 (0.01)	0.00–0.04	0.50
15	0.06	0.06 (0.03)	0.01–0.11	0.5	0.07	0.07 (0.02)	0.02–0.11	0.29	0.06	0.06 (0.02)	0.02–0.10	0.33
16	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.01	0.01 (0.01)	0.00–0.04	1.0	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.04	1.00
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.01	0.02 (0.01)	0.00–0.04	0.5	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
27	0.01	0.01 (0.01)	0.00–0.04	1.0	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.22 Grey Phalarope

Table 28: Grey Phalarope - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.23 Guillemot

Table 29: Guillemot - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.58	0.58 (0.09)	0.42–0.76	0.16	0.63	0.63 (0.10)	0.44–0.83	0.16	0.55	0.56 (0.09)	0.40–0.74	0.16
2	0.95	0.95 (0.17)	0.66–1.31	0.18	0.84	0.84 (0.12)	0.60–1.10	0.14	0.78	0.79 (0.10)	0.60–0.98	0.13
3	0.79	0.79 (0.13)	0.54–1.07	0.16	0.64	0.64 (0.09)	0.46–0.83	0.14	0.60	0.60 (0.08)	0.44–0.77	0.13
4	0.60	0.60 (0.11)	0.39–0.83	0.18	0.55	0.55 (0.08)	0.38–0.70	0.15	0.50	0.50 (0.07)	0.36–0.64	0.14
5	0.36	0.36 (0.08)	0.21–0.51	0.22	0.34	0.34 (0.06)	0.23–0.47	0.18	0.34	0.34 (0.05)	0.23–0.44	0.15
6	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
7	0.01	0.01 (0.01)	0.00–0.04	1.00	0.02	0.02 (0.01)	0.00–0.05	0.50	0.03	0.03 (0.01)	0.01–0.06	0.33
8	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.01 (0.01)	0.00–0.04	1.00
9	0.14	0.14 (0.06)	0.04–0.27	0.43	0.19	0.19 (0.07)	0.08–0.34	0.37	0.22	0.22 (0.06)	0.11–0.35	0.27
10	0.11	0.11 (0.06)	0.01–0.26	0.55	0.10	0.09 (0.04)	0.02–0.18	0.44	0.09	0.09 (0.04)	0.03–0.17	0.44
11	0.04	0.04 (0.02)	0.00–0.09	0.50	0.07	0.07 (0.03)	0.03–0.14	0.43	0.06	0.06 (0.02)	0.02–0.10	0.33
12	0.03	0.03 (0.02)	0.00–0.07	0.67	0.03	0.03 (0.02)	0.00–0.08	0.67	0.03	0.03 (0.02)	0.01–0.08	0.67
13	1.57	1.57 (0.29)	1.04–2.16	0.18	1.33	1.34 (0.21)	0.97–1.79	0.16	1.36	1.35 (0.26)	0.88–1.93	0.19
14	0.43	0.43 (0.12)	0.22–0.68	0.28	0.44	0.44 (0.09)	0.27–0.63	0.20	0.42	0.42 (0.08)	0.27–0.59	0.19
15	0.15	0.15 (0.06)	0.06–0.26	0.40	0.14	0.14 (0.05)	0.06–0.24	0.36	0.19	0.19 (0.05)	0.10–0.29	0.26
16	0.15	0.15 (0.04)	0.08–0.24	0.27	0.11	0.11 (0.03)	0.06–0.18	0.27	0.11	0.11 (0.03)	0.06–0.17	0.27



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.04	0.04 (0.03)	0.00–0.11	0.75	0.03	0.03 (0.02)	0.00–0.07	0.67	0.03	0.03 (0.02)	0.00–0.07	0.67
19	0.26	0.27 (0.07)	0.14–0.40	0.26	0.39	0.39 (0.08)	0.24–0.56	0.21	0.40	0.40 (0.09)	0.23–0.61	0.22
20	0.67	0.66 (0.30)	0.22–1.35	0.45	0.55	0.55 (0.21)	0.24–1.01	0.38	0.54	0.54 (0.17)	0.26–0.91	0.31
21	0.08	0.08 (0.05)	0.00–0.18	0.62	0.11	0.11 (0.04)	0.04–0.19	0.36	0.17	0.17 (0.05)	0.08–0.26	0.29
22	0.24	0.23 (0.07)	0.11–0.39	0.30	0.21	0.21 (0.06)	0.10–0.31	0.29	0.21	0.21 (0.05)	0.12–0.32	0.24
23	0.14	0.14 (0.04)	0.07–0.22	0.29	0.34	0.34 (0.07)	0.21–0.47	0.21	0.39	0.39 (0.07)	0.26–0.54	0.18
24	0.79	0.79 (0.17)	0.48–1.15	0.22	0.66	0.66 (0.12)	0.45–0.89	0.18	0.57	0.56 (0.10)	0.38–0.78	0.18
25	0.84	0.83 (0.14)	0.57–1.13	0.17	0.81	0.81 (0.13)	0.56–1.06	0.16	0.73	0.73 (0.12)	0.50–0.95	0.16
26	0.77	0.77 (0.16)	0.48–1.10	0.21	0.75	0.75 (0.10)	0.56–0.94	0.13	0.73	0.73 (0.10)	0.54–0.92	0.14
27	0.88	0.88 (0.13)	0.62–1.13	0.15	0.77	0.77 (0.11)	0.56–1.00	0.14	0.85	0.84 (0.12)	0.62–1.11	0.14
28	0.50	0.50 (0.15)	0.26–0.81	0.30	0.47	0.46 (0.13)	0.25–0.75	0.28	0.44	0.44 (0.11)	0.26–0.67	0.25
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
30	0.04	0.04 (0.02)	0.00–0.10	0.50	0.04	0.04 (0.02)	0.00–0.09	0.50	0.04	0.04 (0.02)	0.01–0.08	0.50



Table 30: Guillemot Apportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.75	0.76 (0.10)	0.58–0.94	0.13	0.81	0.81 (0.10)	0.60–1.01	0.12	0.72	0.73 (0.09)	0.56–0.91	0.12
2	0.98	0.98 (0.17)	0.69–1.34	0.17	0.88	0.88 (0.13)	0.65–1.15	0.15	0.83	0.84 (0.10)	0.65–1.05	0.12
3	0.79	0.79 (0.13)	0.54–1.07	0.16	0.64	0.64 (0.09)	0.46–0.83	0.14	0.60	0.60 (0.08)	0.44–0.77	0.13
4	0.60	0.60 (0.11)	0.39–0.83	0.18	0.56	0.55 (0.08)	0.39–0.72	0.15	0.51	0.51 (0.07)	0.36–0.65	0.14
5	0.40	0.40 (0.08)	0.23–0.56	0.20	0.37	0.38 (0.06)	0.26–0.51	0.16	0.37	0.37 (0.06)	0.26–0.47	0.16
6	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
7	0.19	0.19 (0.16)	0.00–0.53	0.84	0.14	0.14 (0.12)	0.01–0.45	0.86	0.13	0.13 (0.09)	0.02–0.32	0.69
8	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.01 (0.01)	0.00–0.04	1.00
9	0.61	0.61 (0.13)	0.37–0.85	0.21	0.61	0.63 (0.11)	0.42–0.88	0.17	0.61	0.62 (0.10)	0.42–0.80	0.16
10	0.23	0.23 (0.07)	0.11–0.38	0.30	0.21	0.21 (0.05)	0.11–0.33	0.24	0.26	0.26 (0.06)	0.15–0.38	0.23
11	0.08	0.08 (0.03)	0.02–0.16	0.38	0.11	0.14 (0.04)	0.07–0.22	0.29	0.11	0.11 (0.03)	0.05–0.17	0.27
12	0.06	0.06 (0.02)	0.02–0.11	0.33	0.06	0.06 (0.02)	0.03–0.11	0.33	0.07	0.07 (0.02)	0.03–0.11	0.29
13	1.73	1.73 (0.30)	1.20–2.35	0.17	1.51	1.52 (0.22)	1.12–1.97	0.14	1.60	1.58 (0.27)	1.09–2.18	0.17
14	0.43	0.43 (0.12)	0.22–0.68	0.28	0.45	0.45 (0.09)	0.28–0.65	0.20	0.43	0.43 (0.08)	0.28–0.60	0.19
15	0.17	0.17 (0.06)	0.06–0.28	0.35	0.16	0.16 (0.05)	0.07–0.26	0.31	0.20	0.20 (0.05)	0.11–0.31	0.25
16	0.15	0.15 (0.04)	0.08–0.24	0.27	0.11	0.11 (0.03)	0.06–0.18	0.27	0.11	0.11 (0.03)	0.06–0.17	0.27
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.15	0.15 (0.08)	0.02–0.35	0.53	0.16	0.16 (0.08)	0.04–0.33	0.50	0.14	0.14 (0.06)	0.03–0.27	0.43



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0.29	0.29 (0.07)	0.16–0.43	0.24	0.43	0.42 (0.09)	0.27–0.60	0.21	0.45	0.45 (0.10)	0.28–0.66	0.22
20	0.74	0.73 (0.30)	0.29–1.42	0.41	0.61	0.60 (0.21)	0.29–1.07	0.35	0.63	0.63 (0.18)	0.34–1.01	0.29
21	0.10	0.10 (0.05)	0.01–0.19	0.50	0.12	0.12 (0.04)	0.04–0.21	0.33	0.21	0.20 (0.05)	0.11–0.31	0.25
22	0.24	0.23 (0.07)	0.11–0.39	0.30	0.21	0.21 (0.06)	0.10–0.31	0.29	0.21	0.21 (0.05)	0.12–0.32	0.24
23	0.15	0.15 (0.04)	0.07–0.25	0.27	0.35	0.34 (0.07)	0.21–0.48	0.21	0.40	0.40 (0.07)	0.27–0.56	0.18
24	0.84	0.84 (0.17)	0.53–1.21	0.20	0.72	0.72 (0.12)	0.49–0.96	0.17	0.61	0.61 (0.10)	0.42–0.82	0.16
25	0.84	0.83 (0.14)	0.57–1.13	0.17	0.81	0.81 (0.13)	0.56–1.06	0.16	0.73	0.73 (0.12)	0.50–0.95	0.16
26	0.78	0.78 (0.16)	0.48–1.11	0.21	0.76	0.75 (0.10)	0.56–0.95	0.13	0.74	0.74 (0.10)	0.56–0.93	0.14
27	0.89	0.89 (0.13)	0.64–1.14	0.15	0.78	0.78 (0.11)	0.57–1.00	0.14	0.86	0.86 (0.12)	0.63–1.13	0.14
28	0.51	0.52 (0.15)	0.27–0.82	0.29	0.47	0.47 (0.13)	0.25–0.76	0.28	0.44	0.45 (0.11)	0.26–0.68	0.24
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
30	0.04	0.04 (0.02)	0.00–0.10	0.50	0.04	0.04 (0.02)	0.00–0.09	0.50	0.04	0.04 (0.02)	0.01–0.08	0.50



Table 31: GuillemotApportionedCorrected - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.75	0.76 (0.10)	0.58–0.94	0.13	0.81	0.81 (0.10)	0.60–1.01	0.12	0.72	0.73 (0.09)	0.56–0.91	0.12
2	0.98	0.98 (0.17)	0.69–1.34	0.17	0.88	0.88 (0.13)	0.65–1.15	0.15	0.83	0.84 (0.10)	0.65–1.05	0.12
3	0.79	0.79 (0.13)	0.54–1.07	0.16	0.64	0.64 (0.09)	0.46–0.83	0.14	0.60	0.60 (0.08)	0.44–0.77	0.13
4	0.60	0.60 (0.11)	0.39–0.83	0.18	0.56	0.55 (0.08)	0.39–0.72	0.15	0.51	0.51 (0.07)	0.36–0.65	0.14
5	0.40	0.40 (0.08)	0.23–0.56	0.20	0.37	0.38 (0.06)	0.26–0.51	0.16	0.37	0.37 (0.06)	0.26–0.47	0.16
6	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
7	0.19	0.19 (0.16)	0.00–0.53	0.84	0.14	0.14 (0.12)	0.01–0.45	0.86	0.13	0.13 (0.09)	0.02–0.32	0.69
8	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.01 (0.01)	0.00–0.04	1.00
9	0.61	0.61 (0.13)	0.37–0.85	0.21	0.61	0.63 (0.11)	0.42–0.88	0.17	0.61	0.62 (0.10)	0.42–0.80	0.16
10	0.23	0.23 (0.07)	0.11–0.38	0.30	0.21	0.21 (0.05)	0.11–0.33	0.24	0.26	0.26 (0.06)	0.15–0.38	0.23
11	0.08	0.08 (0.03)	0.02–0.16	0.38	0.11	0.14 (0.04)	0.07–0.22	0.29	0.11	0.11 (0.03)	0.05–0.17	0.27
12	0.06	0.06 (0.02)	0.02–0.11	0.33	0.06	0.06 (0.02)	0.03–0.11	0.33	0.07	0.07 (0.02)	0.03–0.11	0.29
13	1.73	1.73 (0.30)	1.20–2.35	0.17	1.51	1.52 (0.22)	1.12–1.97	0.14	1.60	1.58 (0.27)	1.09–2.18	0.17
14	0.43	0.43 (0.12)	0.22–0.68	0.28	0.45	0.45 (0.09)	0.28–0.65	0.20	0.43	0.43 (0.08)	0.28–0.60	0.19
15	0.17	0.17 (0.06)	0.06–0.28	0.35	0.16	0.16 (0.05)	0.07–0.26	0.31	0.20	0.20 (0.05)	0.11–0.31	0.25
16	0.15	0.15 (0.04)	0.08–0.24	0.27	0.11	0.11 (0.03)	0.06–0.18	0.27	0.11	0.11 (0.03)	0.06–0.17	0.27
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.15	0.15 (0.08)	0.02–0.35	0.53	0.16	0.16 (0.08)	0.04–0.33	0.50	0.14	0.14 (0.06)	0.03–0.27	0.43
19	0.29	0.29 (0.07)	0.16–0.43	0.24	0.43	0.42 (0.09)	0.27–0.60	0.21	0.45	0.45 (0.10)	0.28–0.66	0.22
20	0.74	0.73 (0.30)	0.29–1.42	0.41	0.61	0.60 (0.21)	0.29–1.07	0.35	0.63	0.63 (0.18)	0.34–1.01	0.29
21	0.10	0.10 (0.05)	0.01–0.19	0.50	0.12	0.12 (0.04)	0.04–0.21	0.33	0.21	0.20 (0.05)	0.11–0.31	0.25
22	0.24	0.23 (0.07)	0.11–0.39	0.30	0.21	0.21 (0.06)	0.10–0.31	0.29	0.21	0.21 (0.05)	0.12–0.32	0.24
23	0.15	0.15 (0.04)	0.07–0.25	0.27	0.35	0.34 (0.07)	0.21–0.48	0.21	0.40	0.40 (0.07)	0.27–0.56	0.18
24	0.84	0.84 (0.17)	0.53–1.21	0.20	0.72	0.72 (0.12)	0.49–0.96	0.17	0.61	0.61 (0.10)	0.42–0.82	0.16
25	0.84	0.83 (0.14)	0.57–1.13	0.17	0.81	0.81 (0.13)	0.56–1.06	0.16	0.73	0.73 (0.12)	0.50–0.95	0.16
26	0.78	0.78 (0.16)	0.48–1.11	0.21	0.76	0.75 (0.10)	0.56–0.95	0.13	0.74	0.74 (0.10)	0.56–0.93	0.14
27	0.89	0.89 (0.13)	0.64–1.14	0.15	0.78	0.78 (0.11)	0.57–1.00	0.14	0.86	0.86 (0.12)	0.63–1.13	0.14
28	0.51	0.52 (0.15)	0.27–0.82	0.29	0.47	0.47 (0.13)	0.25–0.76	0.28	0.44	0.45 (0.11)	0.26–0.68	0.24
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
30	0.04	0.04 (0.02)	0.00–0.10	0.50	0.04	0.04 (0.02)	0.00–0.09	0.50	0.04	0.04 (0.02)	0.01–0.08	0.50



2.24 Guillemot/Razorbill

Table 32: Guillemot/Razorbill - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.22	0.22 (0.04)	0.15–0.29	0.18	0.23	0.23 (0.04)	0.16–0.29	0.17	0.22	0.22 (0.03)	0.17–0.29	0.14
2	0.03	0.03 (0.02)	0.00–0.07	0.67	0.05	0.05 (0.02)	0.01–0.09	0.40	0.05	0.05 (0.02)	0.02–0.09	0.40
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
5	0.04	0.04 (0.02)	0.00–0.10	0.50	0.04	0.04 (0.02)	0.01–0.09	0.50	0.03	0.03 (0.02)	0.01–0.06	0.67
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.22	0.22 (0.20)	0.00–0.64	0.91	0.15	0.15 (0.15)	0.00–0.56	1.00	0.12	0.12 (0.11)	0.00–0.35	0.92
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.76	0.75 (0.17)	0.41–1.10	0.23	0.70	0.71 (0.14)	0.44–0.98	0.20	0.62	0.63 (0.12)	0.39–0.88	0.19
10	0.13	0.13 (0.04)	0.06–0.21	0.31	0.13	0.13 (0.04)	0.05–0.21	0.31	0.18	0.18 (0.05)	0.09–0.30	0.28
11	0.05	0.05 (0.03)	0.00–0.11	0.60	0.07	0.07 (0.03)	0.02–0.14	0.43	0.06	0.06 (0.03)	0.02–0.11	0.50
12	0.06	0.06 (0.02)	0.03–0.09	0.33	0.05	0.05 (0.02)	0.02–0.09	0.40	0.06	0.06 (0.02)	0.03–0.09	0.33
13	0.20	0.20 (0.06)	0.09–0.33	0.30	0.22	0.22 (0.07)	0.09–0.37	0.32	0.29	0.29 (0.09)	0.14–0.49	0.31
14	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
15	0.01	0.01 (0.01)	0.00–0.04	1.00	0.02	0.02 (0.01)	0.00–0.05	0.50	0.01	0.02 (0.01)	0.00–0.04	0.50
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.15	0.15 (0.10)	0.00–0.38	0.67	0.18	0.17 (0.10)	0.02–0.39	0.59	0.14	0.14 (0.08)	0.01–0.29	0.57
19	0.05	0.05 (0.03)	0.00–0.10	0.60	0.07	0.07 (0.03)	0.02–0.12	0.43	0.10	0.10 (0.03)	0.04–0.17	0.30
20	0.08	0.08 (0.04)	0.01–0.17	0.50	0.07	0.07 (0.03)	0.01–0.13	0.43	0.10	0.11 (0.05)	0.02–0.23	0.45
21	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.50	0.06	0.06 (0.03)	0.01–0.12	0.50
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.02 (0.01)	0.00–0.04	0.50
24	0.08	0.08 (0.04)	0.01–0.17	0.50	0.10	0.10 (0.03)	0.04–0.16	0.30	0.08	0.08 (0.03)	0.02–0.14	0.38
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
27	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
28	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.25 Gull species

Table 33: Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



2.26 Herring Gull

Table 34: Herring Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
6	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.04	0.04 (0.03)	0.00–0.10	0.75	0.13	0.13 (0.06)	0.03–0.26	0.46	0.16	0.16 (0.05)	0.08–0.27	0.31
10	0.31	0.31 (0.07)	0.17–0.45	0.23	0.30	0.30 (0.06)	0.18–0.40	0.20	0.28	0.28 (0.05)	0.18–0.39	0.18
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.03	0.03 (0.02)	0.00–0.06	0.67	0.03	0.03 (0.02)	0.01–0.07	0.67	0.03	0.03 (0.01)	0.01–0.05	0.33
20	0.10	0.10 (0.03)	0.04–0.14	0.30	0.08	0.08 (0.03)	0.03–0.12	0.38	0.07	0.08 (0.02)	0.04–0.12	0.25
21	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
22	0.06	0.06 (0.05)	0.00–0.17	0.83	0.04	0.04 (0.03)	0.00–0.11	0.75	0.03	0.03 (0.03)	0.00–0.11	1.00
23	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Table 35: HerringGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
6	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.05	0.05 (0.03)	0.00–0.11	0.60	0.13	0.13 (0.06)	0.04–0.27	0.46	0.17	0.17 (0.05)	0.08–0.29	0.29
10	0.32	0.32 (0.07)	0.18–0.46	0.22	0.30	0.31 (0.06)	0.18–0.41	0.19	0.28	0.28 (0.05)	0.18–0.39	0.18
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.03	0.03 (0.02)	0.00–0.06	0.67	0.03	0.03 (0.02)	0.01–0.07	0.67	0.03	0.03 (0.01)	0.01–0.05	0.33
20	0.10	0.10 (0.03)	0.04–0.14	0.30	0.08	0.08 (0.03)	0.03–0.12	0.38	0.07	0.08 (0.02)	0.04–0.12	0.25
21	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
22	0.06	0.06 (0.05)	0.00–0.17	0.83	0.04	0.04 (0.03)	0.00–0.11	0.75	0.03	0.03 (0.03)	0.00–0.11	1.00
23	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.27 Kittiwake

Table 36: Kittiwake - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.46	0.47 (0.09)	0.29–0.66	0.19	0.47	0.48 (0.08)	0.33–0.64	0.17	0.48	0.48 (0.07)	0.34–0.62	0.15
2	0.25	0.25 (0.08)	0.11–0.44	0.32	0.25	0.26 (0.07)	0.15–0.40	0.27	0.31	0.32 (0.12)	0.14–0.57	0.38
3	1.41	1.41 (0.20)	1.04–1.81	0.14	1.58	1.58 (0.31)	1.11–2.24	0.20	1.41	1.43 (0.25)	1.03–2.00	0.17
4	0.51	0.51 (0.16)	0.24–0.86	0.31	0.67	0.67 (0.16)	0.38–1.01	0.24	0.76	0.75 (0.16)	0.46–1.08	0.21
5	0.58	0.57 (0.10)	0.39–0.80	0.18	0.71	0.71 (0.10)	0.52–0.89	0.14	0.65	0.66 (0.09)	0.50–0.84	0.14
6	0.24	0.24 (0.06)	0.13–0.36	0.25	0.36	0.36 (0.12)	0.17–0.63	0.33	0.37	0.37 (0.09)	0.23–0.58	0.24
7	0.55	0.56 (0.17)	0.29–0.90	0.30	0.40	0.40 (0.12)	0.20–0.67	0.30	0.34	0.34 (0.09)	0.19–0.54	0.26
8	1.20	1.20 (0.18)	0.88–1.57	0.15	1.37	1.37 (0.19)	1.02–1.77	0.14	1.38	1.38 (0.17)	1.09–1.71	0.12
9	4.43	4.41 (0.66)	3.29–5.87	0.15	5.28	5.31 (0.92)	3.76–7.36	0.17	4.98	5.00 (0.76)	3.75–6.73	0.15
10	6.48	6.41 (1.40)	4.18–9.52	0.22	6.30	6.27 (1.17)	4.29–8.69	0.19	7.44	7.49 (1.21)	5.30–9.98	0.16
11	0.47	0.47 (0.09)	0.30–0.66	0.19	0.42	0.42 (0.07)	0.28–0.56	0.17	0.45	0.45 (0.07)	0.32–0.57	0.16
12	0.14	0.14 (0.04)	0.06–0.22	0.29	0.17	0.17 (0.03)	0.11–0.24	0.18	0.17	0.17 (0.03)	0.11–0.23	0.18
13	0.76	0.76 (0.17)	0.47–1.12	0.22	0.68	0.67 (0.12)	0.44–0.92	0.18	0.58	0.58 (0.10)	0.40–0.78	0.17
14	0.52	0.53 (0.14)	0.29–0.81	0.26	2.37	2.35 (1.24)	0.53–4.34	0.53	2.71	2.79 (1.13)	0.71–5.15	0.41
15	2.26	2.27 (0.33)	1.60–2.90	0.15	1.91	1.91 (0.25)	1.45–2.41	0.13	1.60	1.61 (0.23)	1.19–2.06	0.14
16	2.42	2.43 (0.51)	1.57–3.58	0.21	3.39	3.37 (0.81)	2.00–5.10	0.24	2.87	2.86 (0.67)	1.66–4.30	0.23
17	0.04	0.04 (0.02)	0.00–0.08	0.50	0.03	0.03 (0.01)	0.00–0.06	0.33	0.04	0.04 (0.02)	0.01–0.08	0.50



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
19	2.09	2.08 (0.39)	1.36–2.88	0.19	2.90	2.89 (0.46)	2.02–3.73	0.16	2.70	2.69 (0.42)	1.94–3.54	0.16
20	2.60	2.59 (0.35)	1.89–3.26	0.14	2.92	2.92 (0.32)	2.34–3.59	0.11	3.19	3.20 (0.34)	2.62–3.88	0.11
21	1.39	1.40 (0.60)	0.42–2.64	0.43	1.91	1.92 (0.55)	0.85–3.02	0.29	1.97	1.97 (0.50)	1.02–2.96	0.25
22	2.09	2.10 (0.73)	1.09–3.69	0.35	1.71	1.73 (0.51)	1.01–2.92	0.29	1.67	1.66 (0.40)	1.07–2.53	0.24
23	2.54	2.54 (0.68)	1.44–4.15	0.27	2.48	2.47 (0.49)	1.73–3.53	0.20	2.71	2.72 (0.41)	2.03–3.58	0.15
24	0.76	0.76 (0.17)	0.45–1.09	0.22	1.04	1.05 (0.37)	0.47–1.88	0.35	1.03	1.04 (0.33)	0.54–1.78	0.32
25	0.87	0.86 (0.24)	0.47–1.36	0.28	1.08	1.08 (0.23)	0.67–1.57	0.21	1.12	1.12 (0.20)	0.75–1.54	0.18
26	0.65	0.66 (0.11)	0.46–0.87	0.17	0.73	0.73 (0.10)	0.54–0.92	0.14	0.64	0.64 (0.08)	0.50–0.80	0.12
27	1.77	1.78 (0.30)	1.23–2.36	0.17	1.69	1.69 (0.24)	1.27–2.15	0.14	1.52	1.53 (0.21)	1.13–1.97	0.14
28	0.58	0.58 (0.14)	0.33–0.88	0.24	0.70	0.70 (0.13)	0.46–0.98	0.19	0.60	0.59 (0.11)	0.38–0.82	0.19
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.07	0.07 (0.03)	0.03–0.13	0.43	0.06	0.06 (0.02)	0.02–0.11	0.33	0.09	0.09 (0.03)	0.05–0.15	0.33



Table 37: KittiwakeApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.46	0.47 (0.09)	0.29–0.66	0.19	0.47	0.48 (0.08)	0.33–0.64	0.17	0.48	0.48 (0.07)	0.34–0.62	0.15
2	0.25	0.25 (0.08)	0.11–0.44	0.32	0.25	0.26 (0.07)	0.15–0.40	0.27	0.31	0.32 (0.12)	0.14–0.57	0.38
3	1.41	1.41 (0.20)	1.04–1.81	0.14	1.58	1.58 (0.31)	1.11–2.24	0.20	1.41	1.43 (0.25)	1.03–2.00	0.17
4	0.51	0.51 (0.16)	0.24–0.86	0.31	0.68	0.68 (0.16)	0.39–1.02	0.24	0.76	0.76 (0.16)	0.47–1.09	0.21
5	0.58	0.57 (0.10)	0.39–0.80	0.18	0.71	0.71 (0.10)	0.52–0.89	0.14	0.65	0.66 (0.09)	0.50–0.84	0.14
6	0.24	0.24 (0.06)	0.13–0.36	0.25	0.36	0.36 (0.12)	0.17–0.63	0.33	0.37	0.37 (0.09)	0.23–0.58	0.24
7	0.55	0.56 (0.17)	0.29–0.90	0.30	0.40	0.40 (0.12)	0.20–0.67	0.30	0.34	0.34 (0.09)	0.19–0.54	0.26
8	1.20	1.20 (0.18)	0.88–1.57	0.15	1.37	1.37 (0.19)	1.02–1.77	0.14	1.38	1.38 (0.17)	1.09–1.71	0.12
9	4.43	4.41 (0.66)	3.29–5.87	0.15	5.28	5.31 (0.92)	3.76–7.36	0.17	4.98	5.00 (0.76)	3.75–6.73	0.15
10	6.48	6.41 (1.40)	4.18–9.52	0.22	6.30	6.27 (1.17)	4.29–8.69	0.19	7.44	7.49 (1.21)	5.30–9.98	0.16
11	0.47	0.47 (0.09)	0.30–0.66	0.19	0.42	0.42 (0.07)	0.28–0.56	0.17	0.45	0.45 (0.07)	0.32–0.57	0.16
12	0.14	0.14 (0.04)	0.06–0.22	0.29	0.17	0.17 (0.03)	0.11–0.24	0.18	0.17	0.17 (0.03)	0.11–0.23	0.18
13	0.78	0.77 (0.17)	0.47–1.14	0.22	0.69	0.68 (0.12)	0.45–0.94	0.18	0.59	0.59 (0.10)	0.41–0.79	0.17
14	0.52	0.53 (0.14)	0.29–0.81	0.26	2.37	2.35 (1.24)	0.53–4.34	0.53	2.71	2.79 (1.13)	0.71–5.15	0.41
15	2.26	2.27 (0.33)	1.60–2.90	0.15	1.91	1.91 (0.25)	1.45–2.41	0.13	1.60	1.61 (0.23)	1.19–2.06	0.14
16	2.42	2.43 (0.51)	1.57–3.58	0.21	3.39	3.37 (0.81)	2.00–5.10	0.24	2.87	2.86 (0.67)	1.66–4.30	0.23
17	0.04	0.04 (0.02)	0.00–0.08	0.50	0.03	0.03 (0.01)	0.00–0.06	0.33	0.04	0.04 (0.02)	0.01–0.08	0.50



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
19	2.09	2.08 (0.39)	1.36–2.88	0.19	2.90	2.90 (0.46)	2.03–3.73	0.16	2.70	2.70 (0.42)	1.95–3.55	0.16
20	2.60	2.59 (0.35)	1.89–3.26	0.14	2.92	2.92 (0.32)	2.34–3.59	0.11	3.19	3.20 (0.34)	2.62–3.88	0.11
21	1.39	1.40 (0.60)	0.42–2.64	0.43	1.91	1.92 (0.55)	0.85–3.02	0.29	1.97	1.97 (0.50)	1.02–2.96	0.25
22	2.09	2.10 (0.73)	1.09–3.69	0.35	1.71	1.73 (0.51)	1.01–2.92	0.29	1.67	1.66 (0.40)	1.07–2.53	0.24
23	2.54	2.54 (0.68)	1.44–4.15	0.27	2.48	2.47 (0.49)	1.73–3.53	0.20	2.71	2.72 (0.41)	2.03–3.58	0.15
24	0.77	0.77 (0.17)	0.46–1.11	0.22	1.05	1.06 (0.37)	0.49–1.90	0.35	1.04	1.05 (0.33)	0.54–1.80	0.31
25	0.87	0.86 (0.24)	0.47–1.36	0.28	1.08	1.08 (0.23)	0.67–1.57	0.21	1.12	1.12 (0.20)	0.75–1.54	0.18
26	0.65	0.66 (0.11)	0.46–0.87	0.17	0.73	0.73 (0.10)	0.54–0.92	0.14	0.64	0.64 (0.08)	0.50–0.80	0.12
27	1.77	1.78 (0.30)	1.23–2.36	0.17	1.69	1.69 (0.24)	1.27–2.15	0.14	1.52	1.53 (0.21)	1.13–1.97	0.14
28	0.58	0.58 (0.14)	0.33–0.88	0.24	0.70	0.70 (0.13)	0.46–0.98	0.19	0.60	0.59 (0.11)	0.38–0.82	0.19
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.07	0.07 (0.03)	0.03–0.13	0.43	0.06	0.06 (0.02)	0.02–0.11	0.33	0.09	0.09 (0.03)	0.05–0.15	0.33



2.28 Large Gull species

Table 38: Large Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.02	0.02 (0.01)	0.00–0.04	0.5
10	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.03	1.0
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.29 Lesser Black-backed Gull

Table 39: Lesser Black-backed Gull - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Table 40: LesserBlackBackedGullApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
17	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	0	0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



2.30 Little Egret

Table 41: Little Egret - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
2	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
3	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
4	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
5	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
6	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
7	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
8	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
9	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
10	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
11	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
12	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
13	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
14	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
15	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
16	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
18	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
19	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
20	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
21	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
22	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
23	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
24	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
25	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
26	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
27	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
28	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
29	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	
30	0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00		0	0.00 (0.00)	0.00–0.00	



2.31 Manx Shearwater

Table 42: Manx Shearwater - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.31	0.30 (0.08)	0.15–0.46	0.27	0.26	0.26 (0.07)	0.15–0.40	0.27	0.25	0.25 (0.06)	0.13–0.38	0.24
2	0.72	0.74 (0.31)	0.23–1.42	0.42	0.98	0.98 (0.35)	0.40–1.75	0.36	1.00	1.00 (0.29)	0.51–1.61	0.29
3	2.70	2.65 (1.84)	0.57–6.82	0.69	2.35	2.30 (1.26)	0.64–5.23	0.55	2.16	2.12 (1.05)	0.69–4.65	0.50
4	0.63	0.64 (0.33)	0.18–1.39	0.52	0.67	0.67 (0.24)	0.29–1.20	0.36	0.65	0.65 (0.21)	0.33–1.15	0.32
5	2.79	2.80 (1.05)	1.08–5.14	0.38	2.34	2.33 (0.74)	1.15–4.01	0.32	2.68	2.66 (0.72)	1.46–4.26	0.27
6	0.85	0.87 (0.27)	0.41–1.43	0.31	0.79	0.79 (0.19)	0.46–1.18	0.24	0.84	0.84 (0.18)	0.54–1.24	0.21
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
13	0.26	0.26 (0.08)	0.12–0.42	0.31	0.26	0.26 (0.06)	0.15–0.38	0.23	0.27	0.27 (0.06)	0.16–0.39	0.22
14	4.00	3.98 (1.38)	1.73–6.89	0.35	4.85	4.81 (1.45)	2.19–7.71	0.30	4.11	4.09 (1.19)	2.09–6.65	0.29
15	1.10	1.09 (0.33)	0.52–1.81	0.30	1.80	1.84 (0.65)	0.79–3.30	0.35	2.29	2.22 (0.86)	0.88–4.23	0.39
16	2.28	2.30 (1.05)	0.76–4.57	0.46	2.18	2.20 (0.74)	1.11–4.00	0.34	1.78	1.79 (0.57)	0.92–3.04	0.32
17	0.34	0.35 (0.09)	0.19–0.52	0.26	0.30	0.30 (0.07)	0.18–0.45	0.23	0.44	0.44 (0.11)	0.26–0.70	0.25



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.06	0.06 (0.02)	0.03–0.10	0.33
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.17	0.17 (0.07)	0.06–0.32	0.41	0.13	0.13 (0.05)	0.05–0.24	0.38	0.13	0.13 (0.04)	0.06–0.21	0.31
25	1.56	1.54 (0.35)	0.94–2.24	0.23	1.55	1.55 (0.31)	1.00–2.17	0.20	1.44	1.44 (0.26)	0.95–1.95	0.18
26	0.26	0.26 (0.08)	0.11–0.44	0.31	0.55	0.56 (0.26)	0.20–1.13	0.46	0.65	0.66 (0.28)	0.24–1.30	0.42
27	1.33	1.33 (0.51)	0.47–2.46	0.38	1.22	1.22 (0.39)	0.57–2.07	0.32	1.19	1.18 (0.32)	0.65–1.88	0.27
28	1.96	1.95 (0.47)	1.14–2.92	0.24	3.12	3.12 (0.95)	1.62–5.31	0.30	3.03	3.01 (0.90)	1.55–4.97	0.30
29	0.32	0.32 (0.07)	0.19–0.46	0.22	0.34	0.34 (0.06)	0.24–0.46	0.18	0.86	0.86 (0.15)	0.57–1.15	0.17
30	0.58	0.58 (0.17)	0.31–0.95	0.29	0.65	0.65 (0.13)	0.41–0.91	0.20	0.63	0.63 (0.11)	0.44–0.87	0.17



Table 43: ManxShearwaterApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.31	0.30 (0.08)	0.15–0.46	0.27	0.26	0.26 (0.07)	0.15–0.40	0.27	0.25	0.25 (0.06)	0.13–0.38	0.24
2	0.72	0.74 (0.31)	0.23–1.42	0.42	0.98	0.98 (0.35)	0.40–1.75	0.36	1.00	1.00 (0.29)	0.51–1.61	0.29
3	2.71	2.67 (1.84)	0.58–6.82	0.69	2.37	2.32 (1.26)	0.66–5.23	0.54	2.18	2.13 (1.05)	0.70–4.67	0.49
4	0.63	0.64 (0.33)	0.18–1.39	0.52	0.67	0.67 (0.24)	0.29–1.20	0.36	0.65	0.65 (0.21)	0.33–1.15	0.32
5	2.79	2.80 (1.05)	1.08–5.14	0.38	2.34	2.34 (0.74)	1.16–4.02	0.32	2.69	2.67 (0.72)	1.48–4.27	0.27
6	0.85	0.87 (0.27)	0.41–1.43	0.31	0.80	0.80 (0.19)	0.46–1.19	0.24	0.86	0.87 (0.18)	0.56–1.26	0.21
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00
13	0.27	0.27 (0.08)	0.13–0.43	0.30	0.28	0.28 (0.07)	0.16–0.42	0.25	0.28	0.28 (0.06)	0.18–0.41	0.21
14	4.00	3.98 (1.38)	1.73–6.89	0.35	4.85	4.81 (1.45)	2.19–7.71	0.30	4.11	4.09 (1.19)	2.09–6.65	0.29
15	1.10	1.09 (0.33)	0.52–1.81	0.30	1.80	1.84 (0.65)	0.79–3.30	0.35	2.29	2.22 (0.86)	0.88–4.23	0.39
16	2.31	2.33 (1.06)	0.80–4.61	0.45	2.20	2.23 (0.74)	1.14–4.02	0.33	1.80	1.81 (0.57)	0.93–3.06	0.31
17	0.37	0.37 (0.09)	0.22–0.55	0.24	0.32	0.31 (0.07)	0.19–0.47	0.23	0.46	0.46 (0.12)	0.28–0.72	0.26



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.00	0.06	0.06 (0.02)	0.03–0.10	0.33
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.02	1.00
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.20	0.20 (0.07)	0.08–0.36	0.35	0.16	0.17 (0.05)	0.07–0.28	0.29	0.15	0.15 (0.04)	0.07–0.24	0.27
25	1.63	1.61 (0.35)	1.00–2.33	0.22	1.60	1.60 (0.32)	1.04–2.23	0.20	1.52	1.52 (0.26)	1.03–2.04	0.17
26	0.26	0.26 (0.08)	0.11–0.44	0.31	0.56	0.57 (0.26)	0.21–1.14	0.46	0.66	0.67 (0.28)	0.24–1.30	0.42
27	1.33	1.33 (0.51)	0.47–2.46	0.38	1.22	1.22 (0.39)	0.57–2.07	0.32	1.19	1.18 (0.32)	0.65–1.88	0.27
28	1.98	1.97 (0.47)	1.15–2.93	0.24	3.13	3.12 (0.95)	1.64–5.32	0.30	3.04	3.01 (0.90)	1.57–4.98	0.30
29	0.32	0.32 (0.07)	0.19–0.46	0.22	0.35	0.35 (0.06)	0.24–0.47	0.17	0.88	0.88 (0.15)	0.60–1.17	0.17
30	0.61	0.61 (0.17)	0.34–0.98	0.28	0.67	0.67 (0.13)	0.43–0.93	0.19	0.64	0.64 (0.11)	0.44–0.90	0.17



2.32 Puffin

Table 44: Puffin - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.06	0.06 (0.03)	0.00–0.13	0.5	0.04	0.04 (0.02)	0.00–0.09	0.50	0.04	0.05 (0.02)	0.01–0.10	0.4
17	0.01	0.01 (0.01)	0.00–0.04	1.0	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.01	0.01 (0.01)	0.00–0.04	1.0	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.05	0.5
28	0.03	0.03 (0.03)	0.00–0.09	1.0	0.03	0.03 (0.02)	0.00–0.08	0.67	0.02	0.02 (0.02)	0.00–0.06	1.0
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Table 45: PuffinApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.06	0.06 (0.03)	0.00–0.13	0.5	0.04	0.04 (0.02)	0.00–0.09	0.50	0.04	0.05 (0.02)	0.01–0.10	0.4
17	0.01	0.01 (0.01)	0.00–0.04	1.0	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.01	0.01 (0.01)	0.00–0.04	1.0	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.05	0.5
28	0.03	0.03 (0.03)	0.00–0.09	1.0	0.03	0.03 (0.02)	0.00–0.08	0.67	0.02	0.02 (0.02)	0.00–0.06	1.0
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Table 46: PuffinApportionedCorrected - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.06	0.06 (0.03)	0.00–0.13	0.5	0.04	0.04 (0.02)	0.00–0.09	0.50	0.04	0.05 (0.02)	0.01–0.10	0.4
17	0.01	0.01 (0.01)	0.00–0.04	1.0	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.01	0.01 (0.01)	0.00–0.04	1.0	0.02	0.02 (0.01)	0.00–0.05	0.50	0.02	0.02 (0.01)	0.00–0.05	0.5
28	0.03	0.03 (0.03)	0.00–0.09	1.0	0.03	0.03 (0.02)	0.00–0.08	0.67	0.02	0.02 (0.02)	0.00–0.06	1.0
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.33 Razorbill

Table 47: Razorbill - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.10	0.10 (0.03)	0.04–0.15	0.30	0.13	0.13 (0.03)	0.07–0.18	0.23	0.12	0.12 (0.03)	0.07–0.17	0.25
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.05	0.05 (0.05)	0.00–0.16	1.00	0.04	0.04 (0.04)	0.00–0.11	1.00	0.03	0.03 (0.03)	0.00–0.09	1.00
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.17	0.17 (0.08)	0.03–0.36	0.47	0.30	0.30 (0.13)	0.10–0.59	0.43	0.28	0.29 (0.11)	0.10–0.55	0.38
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.02)	0.00–0.05	1.00	0.02	0.02 (0.01)	0.00–0.04	0.50
13	0.27	0.28 (0.09)	0.12–0.49	0.32	0.25	0.25 (0.07)	0.13–0.39	0.28	0.25	0.25 (0.06)	0.13–0.38	0.24
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
16	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.40	0.42 (0.39)	0.00–1.21	0.93	0.28	0.28 (0.27)	0.00–0.82	0.96	0.22	0.22 (0.21)	0.00–0.66	0.95
19	0.40	0.40 (0.17)	0.09–0.74	0.42	0.40	0.41 (0.14)	0.15–0.71	0.34	0.43	0.44 (0.13)	0.20–0.71	0.30
20	0.18	0.18 (0.09)	0.03–0.36	0.50	0.17	0.17 (0.06)	0.07–0.31	0.35	0.19	0.19 (0.07)	0.07–0.35	0.37
21	0.21	0.21 (0.10)	0.04–0.42	0.48	0.15	0.15 (0.06)	0.04–0.28	0.40	0.17	0.17 (0.06)	0.06–0.30	0.35
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.49	0.49 (0.11)	0.28–0.73	0.22	0.54	0.54 (0.15)	0.27–0.88	0.28	0.48	0.48 (0.13)	0.26–0.78	0.27
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.09	0.09 (0.05)	0.00–0.20	0.56	0.06	0.06 (0.03)	0.00–0.13	0.50	0.06	0.06 (0.03)	0.01–0.13	0.50
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.21	0.21 (0.11)	0.04–0.46	0.52	0.16	0.16 (0.08)	0.04–0.33	0.50	0.15	0.14 (0.06)	0.05–0.28	0.43
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.16	0.15 (0.12)	0.00–0.43	0.80	0.20	0.21 (0.11)	0.02–0.45	0.52	0.16	0.16 (0.08)	0.02–0.34	0.50



Table 48: RazorbillApportioned - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.20	0.20 (0.04)	0.13–0.27	0.20	0.23	0.23 (0.03)	0.16–0.29	0.13	0.22	0.22 (0.03)	0.16–0.28	0.14
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.10	0.10 (0.07)	0.00–0.25	0.70	0.07	0.07 (0.05)	0.00–0.17	0.71	0.06	0.05 (0.04)	0.00–0.13	0.80
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.46	0.45 (0.11)	0.27–0.67	0.24	0.56	0.57 (0.14)	0.34–0.85	0.25	0.52	0.53 (0.12)	0.33–0.81	0.23
10	0.01	0.01 (0.00)	0.00–0.02	0.00	0.01	0.01 (0.00)	0.00–0.02	0.00	0.02	0.02 (0.00)	0.01–0.03	0.00
11	0.01	0.01 (0.00)	0.00–0.02	0.00	0.01	0.01 (0.00)	0.00–0.02	0.00	0.01	0.01 (0.00)	0.00–0.02	0.00
12	0.05	0.05 (0.02)	0.02–0.10	0.40	0.04	0.04 (0.02)	0.02–0.08	0.50	0.04	0.04 (0.01)	0.02–0.07	0.25
13	0.31	0.32 (0.09)	0.15–0.52	0.28	0.30	0.30 (0.07)	0.16–0.44	0.23	0.31	0.31 (0.06)	0.19–0.45	0.19
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
16	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.44	0.46 (0.39)	0.01–1.26	0.85	0.32	0.32 (0.27)	0.02–0.88	0.84	0.25	0.26 (0.21)	0.01–0.70	0.81



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
19	0.43	0.42 (0.17)	0.11–0.76	0.40	0.44	0.44 (0.14)	0.19–0.75	0.32	0.48	0.49 (0.13)	0.25–0.78	0.27
20	0.20	0.20 (0.09)	0.05–0.38	0.45	0.18	0.18 (0.06)	0.07–0.33	0.33	0.21	0.21 (0.07)	0.09–0.37	0.33
21	0.22	0.22 (0.10)	0.06–0.44	0.45	0.16	0.16 (0.06)	0.05–0.30	0.38	0.20	0.20 (0.06)	0.09–0.33	0.30
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.52	0.52 (0.12)	0.31–0.75	0.23	0.57	0.57 (0.16)	0.30–0.91	0.28	0.50	0.50 (0.13)	0.29–0.80	0.26
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.09	0.09 (0.05)	0.00–0.20	0.56	0.06	0.06 (0.03)	0.00–0.13	0.50	0.06	0.06 (0.03)	0.01–0.13	0.50
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.22	0.21 (0.11)	0.04–0.46	0.52	0.16	0.16 (0.08)	0.04–0.33	0.50	0.15	0.15 (0.06)	0.05–0.28	0.40
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.16	0.15 (0.12)	0.00–0.43	0.80	0.20	0.21 (0.11)	0.02–0.45	0.52	0.16	0.16 (0.08)	0.02–0.34	0.50



Table 49: RazorbillApportionedCorrected - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.20	0.20 (0.04)	0.13–0.27	0.20	0.23	0.23 (0.03)	0.16–0.29	0.13	0.22	0.22 (0.03)	0.16–0.28	0.14
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.10	0.10 (0.07)	0.00–0.25	0.70	0.07	0.07 (0.05)	0.00–0.17	0.71	0.06	0.05 (0.04)	0.00–0.13	0.80
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.46	0.45 (0.11)	0.27–0.67	0.24	0.56	0.57 (0.14)	0.34–0.85	0.25	0.52	0.53 (0.12)	0.33–0.81	0.23
10	0.01	0.01 (0.00)	0.00–0.02	0.00	0.01	0.01 (0.00)	0.00–0.02	0.00	0.02	0.02 (0.00)	0.01–0.03	0.00
11	0.01	0.01 (0.00)	0.00–0.02	0.00	0.01	0.01 (0.00)	0.00–0.02	0.00	0.01	0.01 (0.00)	0.00–0.02	0.00
12	0.05	0.05 (0.02)	0.02–0.10	0.40	0.04	0.04 (0.02)	0.02–0.08	0.50	0.04	0.04 (0.01)	0.02–0.07	0.25
13	0.31	0.32 (0.09)	0.15–0.52	0.28	0.30	0.30 (0.07)	0.16–0.44	0.23	0.31	0.31 (0.06)	0.19–0.45	0.19
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.00
16	0.01	0.01 (0.01)	0.00–0.04	1.00	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.04	1.00
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.44	0.46 (0.39)	0.01–1.26	0.85	0.32	0.32 (0.27)	0.02–0.88	0.84	0.25	0.26 (0.21)	0.01–0.70	0.81
19	0.43	0.42 (0.17)	0.11–0.76	0.40	0.44	0.44 (0.14)	0.19–0.75	0.32	0.48	0.49 (0.13)	0.25–0.78	0.27
20	0.20	0.20 (0.09)	0.05–0.38	0.45	0.18	0.18 (0.06)	0.07–0.33	0.33	0.21	0.21 (0.07)	0.09–0.37	0.33
21	0.22	0.22 (0.10)	0.06–0.44	0.45	0.16	0.16 (0.06)	0.05–0.30	0.38	0.20	0.20 (0.06)	0.09–0.33	0.30
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.52	0.52 (0.12)	0.31–0.75	0.23	0.57	0.57 (0.16)	0.30–0.91	0.28	0.50	0.50 (0.13)	0.29–0.80	0.26
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.09	0.09 (0.05)	0.00–0.20	0.56	0.06	0.06 (0.03)	0.00–0.13	0.50	0.06	0.06 (0.03)	0.01–0.13	0.50
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.22	0.21 (0.11)	0.04–0.46	0.52	0.16	0.16 (0.08)	0.04–0.33	0.50	0.15	0.15 (0.06)	0.05–0.28	0.40
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.16	0.15 (0.12)	0.00–0.43	0.80	0.20	0.21 (0.11)	0.02–0.45	0.52	0.16	0.16 (0.08)	0.02–0.34	0.50



2.34 Red-throated Diver

Table 50: Red-throated Diver - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.01	0.01 (0.01)	0.00–0.05	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.35 Small Gull species

Table 51: Small Gull species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.02	1
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.03	1
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1	0.01	0.01 (0.01)	0.00–0.02	1
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.36 Sooty Shearwater

Table 52: Sooty Shearwater - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.01	0.01 (0.01)	0.00–0.04	1	0.03	0.03 (0.02)	0.00–0.07	0.67	0.02	0.02 (0.01)	0.00–0.05	0.5
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1.00	0.01	0.01 (0.01)	0.00–0.02	1.0
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
30	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



2.37 Unidentified Bird species

Table 53: Unidentified Bird species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.0	0.01	0.01 (0.01)	0.00–0.02	1.0
8	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1.0	0.02	0.02 (0.01)	0.00–0.04	0.5
9	0.01	0.01 (0.01)	0.00–0.04	1	0.01	0.01 (0.01)	0.00–0.03	1.0	0.01	0.01 (0.01)	0.00–0.02	1.0
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
17	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.02	1.0
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.01	0.01 (0.01)	0.00–0.04	1	0.02	0.02 (0.01)	0.00–0.05	0.5	0.02	0.02 (0.01)	0.00–0.04	0.5
30	0.00	0.00 (0.00)	0.00–0.00		0.01	0.01 (0.01)	0.00–0.03	1.0	0.01	0.01 (0.01)	0.00–0.02	1.0



2.38 Wader species

Table 54: Wader species - Densities calculated in 30 digital aerial surveys (DAS) for birds in flight in the Windfarm Development Area (WDA), WDA plus 2 km buffer and WDA plus 4 km buffer. Density is presented with an estimate, mean of bootstraps (Mean) with standard deviation (SD), upper and lower confidence intervals (95% CI) and coefficient of variation (CV).

Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
1	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
2	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
3	0.00	0.00 (0.00)	0.00–0.00		0.02	0.02 (0.02)	0.00–0.06	1.00	0.01	0.02 (0.02)	0.00–0.04	1.00
4	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
5	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
6	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
7	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
8	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
9	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
10	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
11	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
12	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
13	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
14	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
15	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
16	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
17	0.00	0.00 (0.00)	0.00–0.00		0.26	0.26 (0.25)	0.00–0.79	0.96	0.21	0.20 (0.20)	0.00–0.62	1.00



Densities (birds/sq. km.) – Flying												
WDA					WDA + 2 km				WDA + 4 km			
Survey	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV	Estimate	Mean (SD)	95% CI	CV
18	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
19	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
20	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
21	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
22	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
23	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
24	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
25	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
26	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
27	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
28	0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00		0.00	0.00 (0.00)	0.00–0.00	
29	0.09	0.08 (0.08)	0.00–0.26	1.00	0.06	0.06 (0.06)	0.00–0.18	1.00	0.05	0.05 (0.05)	0.00–0.14	1.00
30	0.16	0.16 (0.11)	0.00–0.41	0.69	0.11	0.11 (0.07)	0.00–0.27	0.64	0.08	0.08 (0.06)	0.00–0.21	0.75



